

PROCESSES FOR PRODUCING SUGAR NUCLEOTIDES
AND COMPLEX CARBOHYDRATES

insert #4
H1

TECHNICAL FIELD

This invention relates to a process for producing a complex carbohydrate which is useful for protection against infection of bacteria, viruses and the like, application to cardiovascular disorders and immunotherapy and to a process for producing a sugar nucleotide which is important as a substrate for the synthesis of the complex carbohydrates.

BACKGROUND ART

Examples of the known process for producing sugar nucleotides include: 1) chemical synthetic processes (Adv. Carbohydr. Chem. Biochem., 28, 307 (1973), Bull. Chem. Soc. Japan, 46, 3275 (1973), J. Org. Chem., 57, 146 (1992), Carbohydr. Res., 242, 69 (1993)); 2) production processes using enzymes (J. Org. Chem., 55, 1834 (1990), J. Org. Chem., 57, 152 (1992), J. Am. Chem. Soc., 110, 7159 (1988), Japanese Published Unexamined National Publication No. 508413/95, Japanese Published National Publication No. 500248/95, WO 96/27670); 3) processes using microbial cells such as yeast and the like (Japanese Published Examined Patent Application No. 2073/70, Japanese Published Examined Patent Application No. 40756/71, Japanese Published Examined Patent Application

No. 1837/72, Japanese Published Examined Patent Application No. 26703/72, Japanese Published Examined Patent Application No. 8278/74, Japanese Published Unexamined Patent Application No. 268692/90); and 4) an extraction process from microbial cells of halo-tolerant yeast (Japanese Published Unexamined Patent Application No. 23993/96).

However, the process 1) requires expensive materials (for example, morpholidate derivative of nucleotide-5'-monophosphate (referred to as "NDP" hereinafter), sugar phosphate, etc.); the process 2) requires expensive materials (for example, nucleotide-5'-diphosphate (referred to as "NDP" hereinafter), nucleotide-5'-triphosphate (referred to as "NTP" hereinafter), phosphoenolpyruvate, etc.), and various enzymes (e.g., pyruvate kinase, etc.); and the process 3) requires drying treatment of microbial cells. Including the process 4), all of the above-mentioned processes use expensive nucleotides, sugar phosphates, and the like or have a difficulty in effecting large scale production from the operational point of view, so that an industrial scale production process of sugar nucleotides has not so far been established.

Examples of the known process for producing complex carbohydrates include 1) chemical synthetic processes (Method in *Enzymol.*, 247, 193 (1994), *Angew. Chem. Int. Ed. Engl.*, 21, 155 (1982), *Carbohydr. Res.*, 211, c1 (1991)), 2) processes in

which a hydrolase is used (*Anal. Biochem.*, 202, 215 (1992), *Trends Biotechnol.*, 6, 256 (1988)) and 3) processes in which a glycosyltransferase is used (Japanese Published Unexamined Patent Application No. 79792/95, Japanese Published National Publication No. 500248/95, Japanese Published Examined Patent Application No. 82200/93, WO 94/25614, Japanese Published National Publication No. 503905/97, U.S. Patent 5,583,042).

The introduction of protecting groups is essential for stereo-selective synthesis in the process 1). The yield and selectivity are not sufficient in the process 2). Expensive materials (for example, NDP, NTP, phosphoenolpyruvic acid, sugar phosphate, sugar nucleotide, etc.) and various enzymes (for example, pyruvate kinase, etc.) are necessary in the process 3). Therefore, these processes have not been established as inexpensive industrial production processes of complex carbohydrates. In addition, there has been nothing known about a process for the direct industrial production of complex carbohydrates, which uses only inexpensive nucleotide precursors, sugars and complex carbohydrate precursors as the starting materials.

It has been reported that UMP is produced in a microorganism belonging to the genus *Corynebacterium* when orotic acid is added (*Amino Acid, Nucleic Acid*, 23, 107 (1971)). In addition, a process in which cytidine diphosphate choline is formed from orotic acid as the

material is also known (Japanese Published Unexamined Patent Application No. 276974/93).

DISCLOSURE OF THE INVENTION

An object of the present invention is to provide a process for producing a complex carbohydrate which is useful for protection against infection of bacteria, viruses, and the like, application to cardiovascular disorders and immunotherapy, and a process for producing a sugar nucleotide which is important as a substrate for synthesizing the complex carbohydrate at a low cost and efficiently.

The inventors of the present invention have conducted intensive studies on the microbial production of a complex carbohydrate and a sugar nucleotide using a nucleotide precursor as the starting material, and have found as the results that a sugar nucleotide can be produced by using only a nucleotide precursor and a sugar as the materials, that productivity of the sugar nucleotide can be improved by strengthening expression of genes involved in its biosynthesis and that a complex carbohydrate can be produced efficiently, using a microorganism capable of producing the sugar nucleotide and a microorganism, an animal cell or an insect cell capable of producing the complex carbohydrate from a sugar nucleotide and a complex carbohydrate precursor and using a nucleotide precursor, a sugar and a complex

carbohydrate precursor as the only starting materials, thereby resulting in the accomplishment of the present invention.

The present invention provides a process for producing a complex carbohydrate, which comprises: selecting, as enzyme sources, a) a culture broth of a microorganism capable of producing NTP from a nucleotide precursor, or a treated product of the culture broth, b) a culture broth of a microorganism capable of producing a sugar nucleotide from a sugar and NTP, or a treated product of the culture broth, and c) a culture broth of a microorganism, an animal cell or an insect cell capable of producing a complex carbohydrate from a sugar nucleotide and a complex carbohydrate precursor, or a treated product of the culture broth; allowing the enzyme sources, the nucleotide precursor, the sugar and the complex carbohydrate precursor to be present in an aqueous medium to form and accumulate the complex carbohydrate in the aqueous medium; and recovering the complex carbohydrate from the aqueous medium, a process for producing a sugar nucleotide, which comprises: selecting, as enzyme sources, a) a culture broth of a microorganism capable of producing NTP from a nucleotide precursor, or a treated product of the culture broth, and b) a culture broth of a microorganism capable of producing a sugar nucleotide from a sugar and NTP, or a treated product of the culture broth; allowing the enzyme

sources, the nucleotide precursor and the sugar to be present in an aqueous medium to form and accumulate the sugar nucleotide in the aqueous medium; and recovering the sugar nucleotide from the aqueous medium, and a process for producing a complex carbohydrate, which comprises: selecting, as an enzyme source, a culture broth of a microorganism, an animal cell or an insect cell capable of producing a complex carbohydrate from a sugar nucleotide and a complex carbohydrate precursor, or a treated product of the culture broth; allowing the enzyme source, the sugar nucleotide obtained by the above-described process for producing a sugar nucleotide and the complex carbohydrate precursor to be present in an aqueous medium to form and accumulate the complex carbohydrate in the aqueous medium; and recovering the complex carbohydrate from the aqueous medium. It also provides a process for producing N-acetylglucosamine-1-phosphate, which comprises selecting, as an enzyme source, a culture broth of a microorganism having strong galactokinase activity, or a treated product of the culture broth; allowing the enzyme source and N-acetylglucosamine to be present in an aqueous medium to form and accumulate N-acetylglucosamine-1-phosphate in the aqueous medium; and recovering the N-acetylglucosamine 1-phosphate from the aqueous medium.

BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 shows construction steps of expression plasmids pPA31 and pPAC31.

Fig. 2 shows construction steps of galU, ppa gene expression plasmids pNT12 and pNT32.

Fig. 3 shows construction steps of galT, galK gene expression plasmid pNT25.

Fig. 4 shows construction steps of plasmid pTK7 which expresses galT and galK genes in *Corynebacterium ammoniagenes*.

Fig. 5 shows construction steps of glmU, ppa gene expression plasmid pNT14.

Fig. 6 shows construction steps of pgm gene expression plasmid pNT24.

Fig. 7 shows construction steps of glmM gene expression plasmid pNT44.

Fig. 8 shows construction steps of glk gene expression plasmid pNT46.

Fig. 9 shows construction steps of pfkB gene expression plasmid pNT47.

Fig. 10 shows construction steps of galK gene expression plasmid pNT54.

Fig. 11 shows construction steps of manB, manC gene expression plasmid pNK7.

Fig. 12 shows construction steps of pgm, pfkB gene expression plasmid pNT55.

Fig. 13 shows construction steps of *gmd*, *wcaG* gene expression plasmid pNKG.

Fig. 14 shows construction steps of neuA gene expression plasmid pTA14.

Fig. 15 shows construction steps of lgtC gene expression plasmid pGT3.

Fig. 16 shows construction steps of lgtB gene expression plasmid pNT60.

Abbreviations to be used herein and description of the abbreviations are shown in Table 1-(1) and Table 1-(2).

00000000 001200
00000000 000000

Table 1-(1)

Glc	glucose
G-6-P	glucose-6-phosphate
G-1-P	glucose-1-phosphate
Glc-1,6-P2	glucose-1,6-diphosphate
Gal	galactose
Gal-1-P	galactose-1-phosphate
GlcN-6-P	glucosamine-6-phosphate
GlcN-1-P	glucosamine-1-phosphate
GlcUA	glucuronic acid
GlcN	glucosamine
GlcNAc	N-acetylglucosamine
GlcNAc-1-P	N-acetylglucosamine-1-phosphate
F-6-P	fructose-6-phosphate
F-1,6-P2	fructose-1,6-diphosphate
Man	mannose
Man-6-P	mannose-6-phosphate
Man-1-P	mannose-1-phosphate
GDP-4-keto-6-deoxyMan	guanosine-5'-diphospho- 4-keto-6-deoxymannose
ManNAc	N-acetylmannosamine
NeuAc	N-acetylneuraminic acid
acetyl CoA	acetyl coenzyme A
NTP	nucleotide-5'-triphosphate
NDP	nucleotide-5'-diphosphate
NMP	nucleotide-5'-monophosphate
ATP	adenosine-5'-triphosphate
UTP	uridine-5'-triphosphate
GTP	guanosine-5'-triphosphate
CTP	cytidine-5'-triphosphate
GMP	guanosine-5'-monophosphate

Table 1-(2)

UDP-Glc	uridine-5'-diphosphoglucose
UDP-Gal	uridine-5'-diphosphogalactose
UDP-GlcNAc	uridine-5'-diphospho-N-acetylglucosamine
UDP-GalNAc	uridine-5'-diphospho-N-acetylgalactosamine
UDP-GlcUA	uridine-5'-diphosphoglucuronic acid
GDP-Man	guanosine-5'-diphosphomannose
GDP-Fuc	guanosine-5'-diphosphofucose
CMP-NeuAc	cytidine-5'-monophospho-N-acetylneuraminic acid
galU	glucose-1-phosphate uridylyltransferase
ppa	(inorganic) pyrophosphatase
galK	galactokinase
galT	galactose-1-phosphate uridylyltransferase
glmU	N-acetylglucosamine-1-phosphate uridylyltransferase glucosamine-1-phosphate acetyltransferase
pgm	phosphoglucomutase
pfkB	phosphofructokinase
glmM	phosphoglucosamine mutase
glk	glucokinase
manB	phosphomannomutase
manC	mannose-1-phosphate guanylyltransferase
gmd	GDP-mannose-4,6-dehydratase
wcaG	GDP-4-keto-6-deoxymannose epimerase/reductase
neuA	CMP-N-acetylneuraminic acid synthetase
neuB	N-acetylneuraminic acid synthase
nanA	N-acetylneuraminic acid aldolase
pyrG	cytidine-5'-triphosphate synthetase
lgtB	β 1,4-galactosyltransferase
lgtC	α 1,4-galactosyltransferase
ugd	UDP-glucose dehydrogenase

According to the present invention, a novel production process of a sugar nucleotide and a novel production process of a complex carbohydrate using the sugar nucleotide production process can be provided, which are characterized in that 1) expensive materials (for example, NTP, sugar phosphates, etc.) are not required, and inexpensive nucleotide precursor and a sugar can be used as the sole starting materials, 2) addition of expensive phosphoenolpyruvic acid and pyruvate kinase is not necessary in converting NMP or NDP into NTP, and 3) a process for the isolation of enzymes is not necessary.

With regard to the sugar nucleotide to be produced by the production process of the present invention, compounds having a general structure in which the terminal phosphate group of a nucleoside 5'-diphosphate residue and the reducing group of a sugar residue are linked together by ester bonding can be exemplified, and those compounds in which the nucleotide residue is cytidine 5'-monophosphate and the sugar residue is a polyol are also included in the sugar nucleotide to be produced by the present invention.

Examples of the complex carbohydrate to be produced by the production process of the present invention include compounds in which carbohydrates are bound to monosaccharides, oligosaccharides, monosaccharides or oligosaccharides linked to a carrier or the like, proteins, peptides, lipids,

glycoproteins, glycolipids, glycopeptides, steroid compounds or the like.

The present invention will be described in detail below.

1) With regard to the microorganism for use in the present invention capable of producing NTP from a nucleotide precursor, any microorganism capable of producing NTP from a nucleotide precursor can be used. Examples include microorganisms belonging to the genus *Escherichia* and the genus *Corynebacterium*.

The microorganisms belonging to the genus *Escherichia* include *Escherichia coli* and the like.

The microorganisms belonging to the genus *Corynebacterium* include *Corynebacterium ammoniagenes* and the like.


2) As the microorganism for use in the present invention capable of producing a sugar nucleotide from a sugar and NTP, any microorganism having the activity to form the sugar nucleotide of interest can be used as follows.

2)-(i) With regard to the production of UDP-Glc, it is preferred to use a microorganism having strong enzyme activities of (1) to (4) shown in the following formula 1.

Specifically, microorganisms belonging to the genus *Escherichia* and the genus *Corynebacterium* can be exemplified.

Specific examples include *Escherichia coli* and *Corynebacterium ammoniagenes*.

In addition, a transformant in which at least one of the enzyme activities selected from (1), (2), (3) and (4) are increased by recombinant DNA techniques can also be used. Specific examples of the transformant include *Escherichia coli* KY8415 (FERM BP-408) having recombinant DNA (pNT12) which contains *galU* and *ppa* genes derived from *Escherichia coli*, and the like.



(Formula 1)

- (1): Hexokinase (EC 2.7.1.1) or glucokinase (EC 2.7.1.2)
- (2): Phosphoglucosmutase (EC 2.7.5.1)
- (3): Glucose-1-phosphate uridylyltransferase (EC 2.7.7.9)
- (4): (Inorganic) pyrophosphatase (EC 3.6.1.1)

2)-(ii) With regard to the production of UDP-Gal, it is preferred to use a microorganism having strong enzyme activities of (5) and (6) shown in the following formula 2, preferably further having strong enzyme activities of (1) to (4) shown in the above-mentioned formula 1.

Specifically, microorganisms belonging to the genus *Escherichia* and the genus *Corynebacterium* can be exemplified.

Specific examples include *Escherichia coli* and *Corynebacterium ammoniagenes*.

In addition, a transformant in which at least one of the enzyme activities selected from (5) and (6), or at least one of the enzyme activities selected from (5) and (6) and at least one of the enzyme activities selected from (1) to (4), are increased by recombinant DNA techniques can also be used. Specific examples of the transformant include *Escherichia coli* NM522 having recombinant DNA (pNT25) which contains *galT* and *galK* genes derived from *Escherichia coli* and *Corynebacterium ammoniagenes* ATCC 21170 having recombinant DNA (pTK7) which contains *galT* and *galK* genes derived from *Escherichia coli*.



(Formula 2)

(5): Galactokinase (EC 2.7.1.6)

(6): Galactose-1-phosphate uridylyltransferase (EC 2.7.7.12)

2)-(iii) With regard to the production of UDP-GlcNAc, it is preferred to use a microorganism having strong enzyme activities of (7) to (12) shown in the following formula 3 and having a strong enzyme activity of (4) shown in formula 1, or a microorganism having strong enzyme activities of (13) and (10) shown in formula 3.

Specifically, microorganisms belonging to the genus *Escherichia* and the genus *Corynebacterium* can be exemplified. Specific examples include *Escherichia coli* and *Corynebacterium ammoniagenes*.

In addition, a transformant in which at least one of the enzyme activities selected from (4), (7), (8), (9), (10) and (13) are increased by recombinant DNA techniques can also be used. Specific examples of the transformant include *Escherichia coli* NM522 having recombinant DNA (pNT44) which contains *galM* gene derived from *Escherichia coli*, *Escherichia coli* KY8415 having recombinant DNA (pNT14) which contains *glmU* and *ppa* genes derived from *Escherichia coli*, *Escherichia coli* NM522 having recombinant DNA (pNT46) which contains *glk* gene derived from *Escherichia coli*, *Escherichia coli* NM522 having recombinant DNA (pNT54) which contains *galK* gene derived from *Escherichia coli*, and the like.

Although it is necessary to add Glc-1,6-P2 for the expression and increment of the phosphoglucosamine mutase activity of (8) (*J. Biol. Chem.*, 271, 32 (1996)), it is possible to provide Glc-1,6-P2 from G-6-P and F-6-P without adding Glc-1,6-P2 by using a transformant in which the enzyme activities of (11) and (12) are increased by recombinant DNA techniques.


Specific examples of such a transformant include *Escherichia coli* NM522 having recombinant DNA (pNT24) which

contains *pgm* gene derived from *Escherichia coli*, *Escherichia coli* NM522 having recombinant DNA (pNT47) which contains *pfkB* gene derived from *Escherichia coli*, *Escherichia coli* NM522 having recombinant DNA (pNT55) which contains *pgm* and *pfkB* gene derived from *Escherichia coli*, and the like.

The process in which expression of the phosphoglucosamine mutase activity of (8) is increased by providing Glc-1,6-P2 from G-6-P and F-6-P using the enzyme activities of (11) and (12) is a process disclosed for the first time by the present invention.

The process in which GlcNAc-1-P is produced from GlcNAc using the galactokinase (EC 2.7.1.6) of (13) is a process disclosed for the first time by the present invention. It is possible to produce GlcNAc-1-P by using the process. That is, GlcNAc-1-P can be produced by using a culture broth or a treated product of the culture broth of a microorganism having strong galactokinase activity, such as a microorganism which contains recombinant DNA consisting of a DNA fragment containing a *galK*-encoding gene and a vector, as an enzyme source, allowing the enzyme source and GlcNAc to be present in an aqueous medium to form and accumulate GlcNAc-1-P in the aqueous medium, and recovering the thus-obtained GlcNAc-1-P from the aqueous medium.

Recovery of GlcNAc-1-P from the aqueous medium can be carried out in the usual way in which activated carbon, an ion exchange resin, and the like are used.



(Formula 3)

- (7): Hexokinase (EC 2.7.1.1) or glucokinase (EC 2.7.1.2)
- (8): Phosphoglucosamine mutase
- (9): Glucosamine-1-phosphate acetyltransferase
- (10): N-Acetylglucosamine-1-phosphate uridyltransferase (EC 2.7.7.23)
- (11): Phosphofructokinase (EC 2.7.1.11)
- (12): Phosphoglucomutase (EC 2.7.5.1)
- (13): Galactokinase (EC 2.7.1.6)

2)-(iv) With regard to the production of UDP-GalNAc, it is preferred to use a microorganism having strong enzyme activities of (7) to (12) shown in formula 3, of (14) shown in formula 4 and of (4) shown in formula 1, or a

microorganism having strong enzyme activities of (10) and (13) shown in formula 4 and of (14) shown in formula 4.

Specifically, microorganisms belonging to the genus *Escherichia* and the genus *Corynebacterium* can be exemplified. Specific examples include *Escherichia coli* and *Corynebacterium ammoniagenes*.

In addition, a transformant in which at least one of the enzyme activities selected from (7) to (14) and (4) are increased by recombinant DNA techniques can also be used.

(14)

UDP-GlcNAc \rightleftharpoons UDP-GalNAc

(Formula 4)


(14): UDP-GlcNAc 4-epimerase (EC 5.1.3.7)

2)-(v) With regard to the production of UDP-GlcUA, it is preferred to use a microorganism having strong enzyme activities of (1) to (4) shown in formula 1 and of (15) shown in formula 5.

Specifically, microorganisms belonging to the genus *Escherichia* and the genus *Corynebacterium* can be exemplified. Specific examples include *Escherichia coli* and *Corynebacterium ammoniagenes*.

In addition, a transformant in which at least one of the enzyme activities selected from (1), (2), (3), (4) and

(15) are increased by recombinant DNA techniques can also be used.



(Formula 5)

(15): UDP-Glc dehydrogenase (EC 1.1.1.22)

2)-(vi) With regard to the production of GDP-Man, it is preferred to use a microorganism having strong enzyme activities of (16) to (18) shown in the following formula 6 and of (11) and (12) shown in formula 3.


Specifically, microorganisms belonging to the genus *Escherichia* and the genus *Corynebacterium* can be exemplified. Specific examples include *Escherichia coli* and *Corynebacterium ammoniagenes*.

In addition, a transformant in which at least one of the enzyme activities selected from (16), (17) and (18) are increased by recombinant DNA techniques can also be used. Specific examples of the transformant include *Escherichia coli* NM522 having recombinant DNA (pNK7) which contains *manB* and *manC* genes derived from *Escherichia coli*, *Escherichia coli* NM522 having recombinant DNA (pNT46) which contains *glk* gene derived from *Escherichia coli*, and the like.

Although it is necessary to add Glc-1,6-P2 for the expression and increment of the phosphomannomutase activity

of (17) by recombinant DNA techniques, it is possible to provide Glc-1,6-P₂ from G-6-P and F-6-P without adding Glc-1,6-P₂ by using a transformant in which the enzyme activities of (11) and (12) are increased by recombinant DNA techniques. Specific examples of such a transformant include *Escherichia coli* NM522 having recombinant DNA (pNT24) which contains pgm gene derived from *Escherichia coli*, *Escherichia coli* NM522 having recombinant DNA (pNT47) which contains pfkB gene derived from *Escherichia coli*, *Escherichia coli* NM522 having recombinant DNA (pNT55) which contains pgm and pfkB genes derived from *Escherichia coli*, and the like.

The process in which expression of the phosphomannomutase activity of (17) is increased by providing Glc-1,6-P₂ from G-6-P and F-6-P using the enzyme activities of (11) and (12) is a process disclosed for the first time by the present invention.



- (16): Hexokinase (EC 2.7.1.1) or glucokinase (EC 2.7.1.2)
(17): Phosphomannomutase (EC 2.7.5.7)
(18): Mannose-1-phosphate guanylttransferase (EC 2.7.7.13)

2)-(vii) With regard to the production of GDP-Fuc, it is preferred to use a microorganism having strong enzyme activities of (19) and (20) shown in the following formula 7, of (16) to (18) shown in formula 6 and of (11) and (12) shown in formula 3.

Specifically, microorganisms belonging to the genus *Escherichia* and the genus *Corynebacterium* can be exemplified. Specific examples include *Escherichia coli* and *Corynebacterium ammoniagenes*.

In addition, a transformant in which at least one of the enzyme activities selected from (16), (17), (18), (19) and (20) are increased by recombinant DNA techniques can also be used. Specific examples of the transformant include *Escherichia coli* NM522 having recombinant DNA (pNK7) which contains *manB* and *manC* genes derived from *Escherichia coli*, *Escherichia coli* NM522 having recombinant DNA (pNK8) which contains *gmd* and *wcaG* genes derived from *Escherichia coli*, *Escherichia coli* NM522 having recombinant DNA (pNT46) which contains *glk* gene derived from *Escherichia coli*, and the like.

Although it is necessary to add Glc-1,6-P₂ for the expression and increment of the phosphomannomutase activity of (17) by recombinant DNA techniques, it is possible to provide Glc-1,6-P₂ from G-6-P and F-6-P without adding Glc-1,6-P₂ by using a transformant in which the enzyme

activities of (11) and (12) are increased by recombinant DNA techniques.

Specific examples of such a transformant include *Escherichia coli* NM522 having recombinant DNA (pNT24) which contains *pgm* gene derived from *Escherichia coli*, *Escherichia coli* NM522 having recombinant DNA (pNT47) which contains *pfkB* gene derived from *Escherichia coli*, *Escherichia coli* NM522 having recombinant DNA (pNT55) which contains *pgm* and *pfkB* genes derived from *Escherichia coli*, and the like.




(19): GDP-Man-4,6-dehydratase (EC 4.2.1.47)

(20): GDP-4-keto-6-deoxymannose epimerase/reductase

2)-(viii) With regard to the production of GDP-NeuAc, it is preferred to use a microorganism having strong enzyme activities of (21), (22) or (23), (24) and (25) shown in the following formula 8.

Specifically, microorganisms belonging to the genus *Escherichia* and the genus *Corynebacterium* can be exemplified. Specific examples include *Escherichia coli* and *Corynebacterium ammoniagenes*.

In addition, a transformant in which at least one of the enzyme activities selected from (21), (22), (23), (24) and (25) are increased by recombinant DNA techniques can also be used. Specific examples of the transformant include *Escherichia coli* C600 having recombinant DNA (pNAL1) which contains nanA gene derived from *Escherichia coli* (Appl. Environ. Microbiol., 51, 562 (1986)), *Escherichia coli* NM522 having recombinant DNA (pTA14) which contains neuA gene derived from *Escherichia coli*, and the like.



(Formula 8)

- (21): GlcNAc 2-epimerase (EC 5.1.3.8)
- (22): NeuAc aldolase (EC 4.1.3.3)
- (23): NeuAc synthetase (EC 4.1.3.19)
- (24): CMP-NeuAc synthetase (EC 2.7.7.43)
- (25): CTP synthetase (EC 6.3.4.2)

When a microorganism has both of the properties of microorganisms described in 1) and the properties of microorganisms described in 2), it is possible to produce a sugar nucleotide from a nucleotide precursor and a sugar using the microorganism.

It is possible to produce UDP-Glc from a UTP precursor such as orotic acid or the like and glucose using a microorganism which has both of the properties of microorganisms described in 1) and the properties of microorganisms described in 2)-(i); UDP-Gal from a UTP precursor such as orotic acid or the like and galactose using a microorganism which has both of the properties of microorganisms described in 1) and the properties of microorganisms described in 2)-(ii); UDP-GlcNAc from a UTP precursor such as orotic acid or the like and glucosamine or N-acetylglucosamine using a microorganism which has both of the properties of microorganisms described in 1) and the properties of microorganisms described in 2)-(iii); UDP-GalNAc from a UTP precursor such as orotic acid or the like and glucosamine or N-acetylglucosamine using a microorganism which has both of the properties of microorganisms described in 1) and the properties of microorganisms described in 2)-(iv); UDP-GlcUA from a UTP precursor such as orotic acid or the like and glucose using a microorganism which has both of the properties of microorganisms described in 1) and the properties of microorganisms described in 2)-(v); GDP-Man from a GTP precursor such as GMP or the like and mannose using a microorganism which has both of the properties of microorganisms described in 1) and the properties of

microorganisms described in 2)-(vi); GDP-Fuc from a GTP precursor such as GMP or the like and mannose using a microorganism which has both of the properties of microorganisms described in 1) and the properties of microorganisms described in 2)-(vii); and CMP-NeuAc from a CTP precursor such as orotic acid or the like and N-acetylglucosamine or N-acetylmannosamine using a microorganism which has both of the properties of microorganisms described in 1) and the properties of microorganisms described in 2)-(viii).

Specific examples of such microorganism include *Corynebacterium ammoniagenes* capable of expressing galT and galK genes derived from *Escherichia coli*.

Unlike the case of the above-mentioned strain, when a single strain has only a part of activities required for producing a sugar nucleotide, the sugar nucleotide can be produced by optionally combining microorganisms having respective activities.

The properties described in 1) are not necessarily owned by a single microorganism, and two or more microorganisms in which the properties described in 1) are independently located can also be used as the microorganism having the properties described in 1). Specifically, a combination of *Escherichia coli* capable of expressing *Escherichia coli*-derived pyrG gene with *Corynebacterium*

ammoniogenes (Japanese Published Unexamined Patent Application No. 276974/93) is exemplified.

In the same manner, the microorganism having the properties described in 2) is not necessarily a single microorganism and the properties can independently be owned by two or more microorganisms. By optionally combining the microorganisms, each sugar nucleotide of interest can be produced.

For example, it is possible to produce UDP-Glc from a UTP precursor such as orotic acid or the like and glucose using a microorganism which has the properties of microorganism described in 1) and at least one microorganism having the properties described in 2)-(i); UDP-Gal from a UTP precursor such as orotic acid or the like and galactose using a microorganism which has the properties of microorganisms described in 1) and at least one microorganism having the properties described in 2)-(ii); UDP-GlcNAc from a UTP precursor such as orotic acid or the like and glucosamine or N-acetylglucosamine using a microorganism which has the properties of microorganisms described in 1) and at least one microorganism having the properties described in 2)-(iii); UDP-GalNAc from a UTP precursor such as orotic acid or the like and glucosamine or N-acetylglucosamine using a microorganism which has the properties of microorganisms described in 1) and at least one microorganism having the

properties described in 2)-(iv); UDP-GlcUA from a UTP precursor such as orotic acid or the like and glucose using a microorganism which has the properties of microorganisms described in 1) and at least one microorganism having the properties described in 2)-(v); GDP-Man from a GTP precursor such as GMP or the like and mannose using a microorganism which has the properties described in 1) and at least one microorganism having the properties described in 2)-(vi); GDP-Fuc from a GTP precursor such as GMP or the like and mannose using a microorganism which has the properties described in 1) and at least one microorganism having the properties described in 2)-(vii); and CMP-NeuAc from a CTP precursor such as orotic acid or the like and N-acetylglucosamine or N-acetylmannosamine using a microorganism which has the properties described in 1) and at least one microorganism having the properties described in 2)-(viii).

As described in the foregoing, recombinant microorganisms can be used in the production of sugar nucleotides, and the genes shown in Table 2 related to the production of sugar nucleotides have been cloned from the chromosome of *Escherichia coli* and their complete nucleotide sequences have been determined.

Table 2

Genes	References
galU gene	<i>J. Biochem.</i> , 115, 965 (1994)
ppa gene	<i>J. Bacteriol.</i> , 170, 5901 (1988)
galK gene	<i>Nucleic Acids Res.</i> , 13, 1841 (1985)
galT gene	<i>Nucleic Acids Res.</i> , 14, 7705 (1986)
glmU gene	<i>J. Bacteriol.</i> , 175, 6150 (1993)
pgm gene	<i>J. Bacteriol.</i> , 176, 5847 (1994)
pfkB gene	<i>Gene</i> , 28, 337 (1984)
glmM gene	<i>J. Biol. Chem.</i> , 271, 32 (1996)
glk gene	<i>J. Bacteriol.</i> , 179, 1298 (1997)
manB gene	<i>J. Bacteriol.</i> , 178, 4885 (1996)
manC gene	<i>J. Bacteriol.</i> , 178, 4885 (1996)
gmd gene	<i>J. Bacteriol.</i> , 178, 4885 (1996)
wcaG gene	<i>J. Bacteriol.</i> , 178, 4885 (1996)
neuA gene	<i>J. Biol. Chem.</i> , 264, 14769 (1989)
neuB gene	<i>J. Bacteriol.</i> , 177, 312 (1995)
nanA gene	<i>Nucleic Acids Res.</i> , 13, 8843 (1985)
pyrG gene	<i>J. Biol. Chem.</i> , 261, 5568 (1986)
ugd gene	<i>J. Bacteriol.</i> , 177, 4562 (1995)

Various procedures related to recombinant DNA techniques, such as isolation and purification of plasmid DNA from *Escherichia coli* having a plasmid which contains the genes, cleavage of the plasmid DNA with restriction enzymes,

isolation and purification of the cleaved DNA fragments, enzymatic linking of the DNA fragments and transformation of *Escherichia coli* using recombinant DNA, can be carried out in accordance with known processes (for example, J. Sambrook et al., Molecular Cloning, A Laboratory Manual, second edition, Cold Spring Harbor Laboratory (1989)). In addition, the polymerase chain reaction (referred to as "PCR" hereinafter) can be carried out, for example, using Thermal Cycler manufactured by Perkin-Elmer-Cetus.

Expression of a gene related to the production of a sugar nucleotide in a host can be effected by a procedure in which a DNA fragment containing the gene is obtained as an appropriate length of DNA fragment containing the gene using restriction enzymes or PCR and then the thus formed DNA fragment is inserted into downstream of the promoter of an expression vector, and the DNA-inserted expression vector is introduced into a host cell which is suited for the expression vector.

Every microorganism can be used as the host, so long as it can express the gene of interest. Examples include microorganisms belonging to the genus *Escherichia*, *Serratia*, *Corynebacterium*, *Brevibacterium*, *Pseudomonas*, *Bacillus* and the like, as well as yeasts belonging to the genus *Saccharomyces*, *Candida* and the like.

With regard to the expression vector to be used, those having an ability to replicate autonomously in the above-described host or to be integrated into chromosome, and containing a promoter at the position at which transcription of the gene related to the production of a sugar nucleotide can be effected, may be used.

When the above-mentioned microorganism is used as the host, it is preferred that the expression vector of a gene related to the production of a sugar nucleotide can be replicated autonomously in the microorganism and that, at the same time, the expression vector comprises a promoter, a ribosome-binding sequence, the gene related to the production of sugar nucleotide and a transcription terminator sequence. It may also contain a regulatory gene of the promoter.

Examples of the expression vector include pBTrp2, pBTac1, pBTac2 (all manufactured by Boehringer Mannheim Co.), pKYP10 (Japanese Published Unexamined Patent Application No. 110600/83), pKYP200 (Agric. Biol. Chem., 48, 669 (1984)), pLSA1 (Agric. Biol. Chem., 53, 277 (1989)), pGEL1 (Proc. Natl. Acad. Sci. USA., 82, 4306 (1985)), pBluescript II SK+ (manufactured by STRATAGENE), pTrs30 (prepared from *Escherichia coli* JM109/pTrs30 (FERM BP-5407)), pTrs32 (prepared from *Escherichia coli* JM109/pTrs32 (FERM BP-5408)), pUC19 (Gene, 33, 103 (1985)), pSTV28 (manufactured by Takara Shuzo Co., Ltd.), pPA1 (Japanese Published Unexamined Patent

Application No. 233798/88), pCG11 (Japanese Examined Patent Application No. 91827/94), and the like.

Any promoter can be used, so long as it can be expressed in the above-mentioned host. Examples include promoters derived from *Escherichia coli*, phage and the like, such as trp promoter, lac promoter, P_L promoter, P_R promoter, and the like. Also usable are artificially designed and modified promoters such as trp tandem promoter in which two trp promoters are connected in series, tac promoter, and the like.

With regard to the ribosome-binding sequence, any sequence capable of being expressed in the above-mentioned host can be used, but it is preferred to use a plasmid in which the region between a ribosome-binding sequence and an initiation codon is adjusted to a suitable distance (for example, 6 to 18 bases).

Although the transcription terminator sequence is not always necessary for the expression of genes related to the production of sugar nucleotide, it is preferred to arrange the transcription termination sequence preferably at a downstream position of the structural gene.

Any microorganism can be used as the host, so long as it can express the recombinant DNA and can apply to the sugar nucleotide formation reaction. Examples include *Escherichia coli* XL1-Blue, *Escherichia coli* XL2-Blue, *Escherichia coli*

[illegible][illegible]

Case

parapsilosis, *Candida krusei*, *Candida versatilis*, *Candida lipolytica*, *Candida zeylanoides*, *Candida guilliermondii*, *Candida albicans*, *Candida humicola*, *Pichia farinosa*, *Pichia ohmeri*, *Torulopsis candida*, *Torulopsis sphaerica*, *Torulopsis xylinus*, *Torulopsis famata*, *Torulopsis versatilis*, *Debaryomyces subglobosus*, *Debaryomyces cantarellii*, *Debaryomyces globosus*, *Debaryomyces hansenii*, *Debaryomyces japonicus*, *Zygosaccharomyces rouxii*, *Zygosaccharomyces bailii*, *Kluyveromyces lactis*, *Kluyveromyces marxianus*, *Hansenula anomala*, *Hansenula jadinii*, *Brettanomyces lambicus*, *Brettanomyces anomalus*, *Schizosaccharomyces pombe*, *Trichosporon pullulans*, *Schwanniomyces alluvius*, and the like.

Culturing of the microorganism for use in the present invention can be carried out in accordance with the usual culturing process. The medium for use in the culturing of the microorganism may be either a nutrient medium or a synthetic medium, so long as it contains carbon sources, nitrogen sources, inorganic salts and the like which can be assimilated by the microorganism and it can perform culturing of the microorganism efficiently.

Examples of the carbon sources include those which can be assimilated by each microorganism, such as carbohydrates (for example, glucose, fructose, sucrose, lactose, maltose, mannitol, sorbitol, molasses, starch, starch hydrolysate, etc.), organic acids (for example,

pyruvic acid, lactic acid, citric acid, fumaric acid, etc.), various amino acids (for example, glutamic acid, methionine, lysine, etc.), and alcohols (for example, ethanol, propanol, glycerol, etc.). Also useful are natural organic nutrient sources, such as rice bran, cassava, bagasse, corn steep liquor, and the like.

Examples of the nitrogen sources include various inorganic and organic ammonium salts (for example, ammonia, ammonium chloride, ammonium sulfate, ammonium carbonate, ammonium acetate, ammonium phosphate, etc.), amino acids (for example, glutamic acid, glutamine, methionine, etc.), peptone, NZ amine, corn steep liquor, meat extract, yeast extract, malt extract, casein hydrolysate, soybean meal, fish meal or a hydrolysate thereof and the like.

Examples of the inorganic substances include potassium dihydrogen phosphate, dipotassium hydrogen phosphate, sodium dihydrogen phosphate, disodium hydrogen phosphate, magnesium phosphate, magnesium sulfate, magnesium chloride, sodium chloride, calcium chloride, ferrous sulfate, manganese sulfate, copper sulfate, zinc sulfate, calcium carbonate, and the like. Vitamins, amino acids, nucleic acids and the like may be added as occasion demands.

The culturing is carried out under aerobic conditions by shaking culture, aeration stirring culture or the like means. The culturing temperature is preferably from 15 to

45°C, and the culturing time is generally from 5 to 96 hours. The pH of the medium is maintained at 3.0 to 9.0 during the culturing. Adjustment of the medium pH is carried out using an inorganic or organic acid, an alkali solution, urea, calcium carbonate, ammonia and the like. Also, antibiotics (for example, ampicillin, tetracycline, etc.) may be added to the medium during the culturing as occasion demands.

When a microorganism transformed with an expression vector in which an inducible promoter is used as the promoter is cultured, an inducer may be added to the medium as occasion demands. For example, isopropyl- β -D-thiogalactopyranoside (IPTG) or the like may be added to the medium when a microorganism transformed with an expression vector containing lac promoter is cultured, or indoleacrylic acid (IAA) or the like may be added thereto when a microorganism transformed with an expression vector containing trp promoter is cultured.

When two or more microorganisms are used in the sugar nucleotide production of the present invention, the microorganisms may be separately cultured to use the resulting culture broths in the sugar nucleotide production or inoculated simultaneously into a single culture vessel to carry out mixed culturing and to use the resulting culture broth in the sugar nucleotide production. In an alternative way, one of the microorganisms is firstly cultured, the

remaining microorganism is inoculated during or after the culturing and cultured, and then the resulting culture broth is used in the sugar nucleotide production. In another alternative way, a microorganism having the properties described in 1) and a microorganism having the properties described in 2) may be separately cultured and used in the sugar nucleotide production using the resulting culture broths.

The microbial culture broth obtained by the culturing or a treated product of the culture broth obtained by treating the culture broth in various ways can be used as an enzyme source for the formation of a sugar nucleotide in an aqueous medium.

Examples of the treated product of the culture broth include a concentrated product of the culture broth, a dried product of the culture broth, a culture supernatant obtained by centrifuging the culture broth, a concentrated product of the culture supernatant, an enzyme preparation obtained from the culture supernatant, cells (including microbial cells) obtained by centrifuging the culture broth, a dried product of the cells, a freeze-dried product of the cells, a surfactant-treated product of the cells, an ultrasonic-treated product of the cells, a mechanically disrupted product of the cells, a solvent-treated product of the cells, an enzyme-treated product of the cells, a protein fraction of

the cells, an immobilized product of the cells and an enzyme preparation obtained by extraction from the cells.

The amount of the enzyme source used in the formation of the sugar nucleotide is within the range of from 1 to 500 g/l, preferably from 5 to 300 g/l, as wet cells. When the reaction is carried out in an aqueous medium using two or more microorganisms simultaneously, amount of the total wet cells of the microorganisms in the aqueous medium is within the range of from 2 to 500 g/l, preferably from 5 to 400 g/l.

Examples of the aqueous medium used in the formation of the sugar nucleotide include water, buffer solutions (for example, those of phosphate, carbonate, acetate, borate, citrate, Tris, etc.), alcohols (for example, methanol, ethanol, etc.), esters (for example, ethyl acetate, etc.), ketones (for example, acetone, etc.), amides (for example, acetamide, etc.), and the like. The microbial culture broth used as the enzyme source may also be used as the aqueous medium.

Examples of the nucleotide precursor used in the formation of the sugar nucleotide include orotic acid, uracil, orotidine, uridine, cytosine, cytidine, adenine, adenosine, guanine, guanosine, hypoxanthine, inosine, xanthine, xanthosine, inosine-5'-monophosphate, xanthosine-5'-monophosphate, guanosine-5'-monophosphate, uridine-5'-monophosphate, cytidine-5'-monophosphate, and the like.

Preferred are orotic acid and guanosine-5'-monophosphate. The nucleotide precursor may be in the form of a purified product or in the form of a salt of the precursor, and a culture broth containing the precursor produced by the fermentation of a microorganism or the precursor roughly purified from the culture broth may also be used as the nucleotide precursor, so long as its impurities do not inhibit the reaction. The nucleotide precursor is used at a concentration of from 0.1 mM to 1.0 M, preferably from 0.01 to 0.3 M.

Examples of the sugar used in the formation of the sugar nucleotide include glucose, fructose, galactose, glucosamine, N-acetylglucosamine, N-acetylgalactosamine, mannose, fucose, N-acetylmannosamine, ^{N-}acetylneuraminic acid and the like, and derivatives thereof. The sugar may be either in the form of a purified product or in the form of a material containing the same, so long as impurities in the material do not inhibit the reaction. The sugar is used at a concentration of from 0.1 mM to 2.0 M, by adding it in one portion when the reaction is started or in portions or continuously during the reaction.

In the formation of the sugar nucleotide, an energy source necessary for the regeneration of ATP, a coenzyme, a phosphate ion, a magnesium ion, a chelating agent (for

example, phytic acid, etc.), a surfactant and an organic solvent may be added as occasion demands.

Examples of the energy source include carbohydrate (for example, glucose, fructose, sucrose, lactose, maltose, mannitol, sorbitol, ^{molasses, starch hydrolysate} etc.), organic acids (for example, pyruvic acid, lactic acid, acetic acid, etc.), amino acids (for example, glycine, alanine, aspartic acid, glutamic acid, etc.), ~~molasses, starch hydrolysate~~, and the like, which may be used at a concentration of from 1.0 mM to 2.0 M.

Examples of the phosphate ion include orthophosphoric acid, polyphosphoric acids (for example, pyrophosphoric acid, tripolyphosphoric acid, tetrapolyphosphoric acid, tetrapoly metaphosphoric acid, etc.), polymetaphosphoric acids, inorganic phosphates (for example, potassium dihydrogen phosphate, dipotassium hydrogen phosphate, sodium dihydrogen phosphate, disodium hydrogen phosphate, etc.), and the like, which may be used at a concentration of from 1.0 mM to 1.0 M.

Examples of the magnesium ion include inorganic magnesium salts (for example, magnesium sulfate, magnesium nitrate, magnesium chloride, etc.), organic magnesium salts (for example, magnesium citrate, etc.), and the like, which may be used at a concentration of generally from 1 to 100 mM.

Examples of the surfactant include those which can enhance the production of various sugar nucleotides, such as nonionic surfactants (for example, polyoxyethylene

octadecylamine (e.g., Nymean S-215, manufactured by Nippon Oils and Fats Co.), etc.), cationic surfactants (for example, cetyl trimethylammonium bromide, alkyltrimethylbenzylammonium chloride (e.g., Cation F2-40E, manufactured by Nippon Oils and Fats Co.) etc.), anionic surfactants (for example, lauroyl sarcosinate, etc.) and tertiary amines (for example, alkyltrimethylamine (e.g., Tertiary Amine FB, manufactured by Nippon Oils and Fats Co.), etc.), which may be used alone or as a mixture of two or more. The surfactant may be used at a concentration of generally from 0.1 to 50 g/l.

Examples of the organic solvent include xylene, toluene, aliphatic alcohol, acetone, ethyl acetate, and the like, which may be used at a concentration of generally from 0.1 to 50 ml/l.

The reaction for forming a sugar nucleotide can be carried out in an aqueous medium at pH of from 5 to 10, preferably from 6 to 9, at a temperature of from 20 to 50°C and for a period of from 2 to 96 hours.

The sugar nucleotide can be formed by the process. Examples include a uridine diphosphate compound, a guanosine diphosphate compound, a cytidine monophosphate compound and the like. Specific examples include sugar nucleotides selected from UDP-Glc, UDP-Gal, UDP-GlcNAc, UDP-GalNAc, UDP-GlcUA, GDP-Man, GDP-Fuc, CMP-NeuAc, and the like, and derivatives thereof.

Determination of the sugar nucleotide formed in the aqueous medium can be carried out in accordance with a known method, for example, isolation and determination of UDP-Glc and UDP-Gal can be carried out by high performance liquid chromatography (referred to as "HPLC" hereinafter) method described in *Anal. Biochem.*, 216, 188 (1994). In addition, isolation and determination of UDP-GlcNAc, GDP-Man, GDP-Fuc and CMP-NeuAc can be carried out by HPLC under the following conditions:

Elution solution:

0.1 M KH_2PO_4 (adjust to pH 3.2 with H_3PO_4)

Flow rate:

1 ml/min

Column:

Partisil-10 SAX (manufactured by Whatman)

Detection:

UV 262 nm

Determination:

Calculated by comparing standard absorbance values

Recovery of the sugar nucleotide formed in the reaction solution can be carried out in the usual way using activated carbon, an ion exchange resin and the like, for example, UDP-Gal and UDP-Glc can be recovered in accordance with the process described in *J. Org. Chem.*, 57, 152 (1992),

and UDP-GlcNAc with the process described in *J. Org. Chem.*,
57, 146 (1992).

With regard to the microorganisms, animal cells or insect cells eligible for in the production of the complex carbohydrate of the present invention, all microorganisms, animal cells or insect cells capable of producing the complex carbohydrate from a sugar nucleotide and a complex carbohydrate precursor can be used. Examples of such microorganisms, animal cells or insect cells include those which have the activities of glucosyltransferase, galactosyltransferase, N-acetylglucosaminyltransferase, N-acetylgalactosaminyltransferase, glucuronosyltransferase, mannosyltransferase, sialyltransferase, fucosyltransferase, and the like.

In addition, microorganisms, animal cells or insect cells modified by recombinant DNA techniques can also be used in a similar way to the case of the above-mentioned sugar nucleotide production. Examples of such microorganisms, animal cells or insect cells include *Escherichia coli* which expresses the ceramide glucosyltransferase gene derived from human melanoma cell line SK-Mel-28 (*Proc. Natl. Acad. Sci. USA.*, 93, 4638 (1996)), human melanoma cell line WM266-4 which produces β 1,3-galactosyltransferase (ATCC CRL 1676), recombinant cell line such as namalwa cell line KJM-1 or the

like which contains the β 1,3-galactosyltransferase gene derived from the human melanoma cell line WM266-4 (Japanese Published Unexamined Patent Application No. 181759/94), *Escherichia coli* (EMBO J., 9, 3171 (1990)) or *Saccharomyces cerevisiae* (Biochem. Biophys. Res. Commun., 201, 160 (1994)) which expresses the β 1,4-galactosyltransferase gene derived from human HeLa cells, COS-7 cell line (ATCC CRL 1651) which expresses the rat β 1,6-N-acetylglucosaminyltransferase gene (J. Biol. Chem., 268, 15381 (1993)), Sf9 cell line which expresses human N-acetylglucosaminyltransferase gene (J. Biochem., 118, 568 (1995)), *Escherichia coli* which expresses human glucuronosyltransferase (Biochem. Biophys. Res. Commun., 196, 473 (1993)), namalwa cell line which expresses human α 1,3-fucosyltransferase (J. Biol. Chem., 269, 14730 (1994)), COS-1 cell line which expresses human α 1,3/1,4-fucosyltransferase (Genes Dev., 4, 1288 (1990)), COS-1 cell line which expresses human α 1,2-fucosyltransferase (Proc. Natl. Acad. Sci. USA., 87, 6674 (1990)), COS-7 cell line which expresses chicken α 2,6-sialyltransferase (Eur. J. Biochem., 219, 375 (1994)), COS cell line which expresses human α 2,8-sialyltransferase (Proc. Natl. Acad. Sci. USA., 91, 7952 (1994)), *Escherichia coli* which expresses β 1,3-N-acetylglucosaminyltransferase, β 1,4-galactosyltransferase, β 1,3-N-acetylgalactosaminyltransferase or α 1,4-galactosyltransferase derived from *Neisseria* (WO 96/10086),

Escherichia coli which expresses *Neisseria*-derived α 2,3-sialyltransferase (*J. Biol. Chem.*, 271, 28271 (1996)), *Escherichia coli* which expresses *Helicobacter pylori*-derived α 1,3-fucosyltransferase (*J. Biol. Chem.*, 272, 21349 and 21357 (1997)), *Escherichia coli* which expresses yeast-derived α 1,2-mannosyltransferase (*J. Org. Chem.*, 58, 3985 (1993)), and the like.

When a microorganism is used for producing the complex carbohydrate of the present invention, the microorganism can be cultured using the same medium under the same culture conditions as in the case of the above-mentioned microorganism capable of producing a sugar nucleotide from a nucleotide precursor and a sugar.

When animal cells are used for producing the complex carbohydrate of the present invention, the preferred culture medium is generally RPMI 1640 medium, Eagle's MEM medium or a medium thereof modified by further adding fetal calf serum, and the like. The culturing is carried out under certain conditions, for example, in the presence of 5% CO₂. The culturing is carried out at a temperature of preferably from 20 to 40°C for a period of generally from 3 to 14 days. As occasion demands, antibiotics may be added to the medium.

When insect cells are used for producing the complex carbohydrate of the present invention, culturing of the

insect cells can be carried out in accordance with the known process (*J. Biol. Chem.*, 268, 12609 (1993)).

The culture broth of a microorganism, an animal cell line or an insect cell line obtained by the culturing and a treated product of the culture broth obtained by treating the culture broth in various ways can be used as an enzyme source for forming the complex carbohydrate in an aqueous medium.

Examples of the treated product of the culture broth include a concentrated product of the culture broth, a dried product of the culture broth, a culture supernatant obtained by centrifuging the culture broth, a concentrated product of the culture supernatant, an enzyme preparation obtained from the culture supernatant, cells (including microbial cells) obtained by centrifuging the culture broth, a dried product of the cells, a freeze-dried product of the cells, a surfactant-treated product of the cells, an ultrasonic-treated product of the cells, a mechanically disrupted product of the cells, a solvent-treated product of the cells, an enzyme-treated product of the cells, a protein fraction of the cells, an immobilized product of the cells and an enzyme preparation obtained by extraction from the cells.

The enzyme source used in the formation of the complex carbohydrate is typically within the range of from 0.1 mU/l to 10,000 U/l, preferably from 1 mU/l to 1,000 U/l (where 1 unit (U) is the amount of the enzyme activity which

can form 1 μ mole of the complex carbohydrate within 1 minute at 37°C).

Examples of the aqueous medium used in the formation of the complex carbohydrate include water, buffer solutions (for example, those of phosphate, carbonate, acetate, borate, citrate, Tris, etc.), alcohols (for example, methanol, ethanol, etc.), esters (for example, ethyl acetate, etc.), ketones (for example, acetone, etc.), amides (for example, acetamide, etc.), and the like. Each of the culture broths of microorganisms, animal cells or insect cells used as the enzyme sources may also be used as the aqueous medium.

As occasion demands, chelating agents (for example, phytic acid, etc.), inorganic salts (for example, $MnCl_2$, etc.), β -mercaptoethanol and the like may be added.

As the sugar nucleotide used in the formation of the complex carbohydrate, the above-mentioned reaction solution obtained by the sugar nucleotide formation or the sugar nucleotide purified from the reaction solution can be used at a concentration of from 0.01 mM to 2.0 M.

In addition, a sugar nucleotide can be supplied in the complex carbohydrate formation reaction solution by forming the sugar nucleotide by the above-mentioned process.

With regard to the complex carbohydrate precursor used in the formation of the complex carbohydrate, any material can be used, so long as it can be used as the

substrate of glycosyltransferase. Examples include monosaccharides, oligosaccharides, monosaccharides or oligosaccharides linked to a carrier or the like, proteins, peptides, lipids, glycoproteins, glycolipids, glycopeptides, steroid compounds, and the like.

Specific examples include glucose, galactose, mannose, sialic acid, N-acetylglucosamine, N-acetylgalactosamine, lactose, N-acetyllactosamine, lacto-N-biose, GlcNAc β 1-3Gal β 1-4Glc, GlcNAc β 1-4Gal β 1-4Glc, globotriose, Gal α 1-4Gal β 1-4GlcNAc, 2'-fucosyllactose, 3-fucosyllactose, 3'-sialyllactose, 6'-sialyllactose, 3'-sialyl-N-acetyllactosamine, 6'-sialyl-N-acetyllactosamine, sialyllacto-N-biose, Lewis X, Lewis a, lacto-N-tetraose, lacto-N-neotetraose, lactodifucotetraose, 3'-sialyl-3-fucosyllactose, sialyl-Lewis X, sialyl-Lewis a, lacto-N-fucopentaose I, lacto-N-fucopentaose II, lacto-N-fucopentaose III, lacto-N-fucopentaose V, LS-tetrasaccharide a, LS-tetrasaccharide b, LS-tetrasaccharide c, ¹(α 2,3)sialyllacto-N-neotetraose and derivatives thereof, serine, threonine, asparagine and peptides containing these amino acids and derivatives thereof, ceramide and derivatives thereof, and the like. The complex carbohydrate precursor can be used at a concentration of from 0.01 mM to 2.0 M.

Examples of the complex carbohydrate of the present invention include complex carbohydrates containing at least one sugar selected from glucose, galactose,

[illegible]

GalNAc β 1-3Gal, GalNAc β 1-4Gal, GalNAc β 1-4GlcNAc,
GalNAc α 1-3GalNAc, Man β 1-4GlcNAc, Man α 1-6Man, Man α 1-3Man,
Man α 1-2Man, GlcUA β 1-4GlcN, GlcUA β 1-3Gal, GlcUA β 1-3GlcNAc,
GlcUA β 1-3GalNAc, NeuAc α 2-3Gal, NeuAc α 2-6Gal, NeuAc α 2-3GlcNAc,
NeuAc α 2-6GlcNAc, NeuAc α 2-3GalNAc, NeuAc α 2-6GalNAc,
NeuAc α 2-8NeuAc, Fuc α 1-3Glc, Fuc α 1-4Glc, Fuc α 1-3GlcNAc,
Fuc α 1-4GlcNAc, Fuc α 1-2Gal and Fuc α 1-6GlcNAc; and complex
carbohydrates which contain the just described complex
carbohydrates. In this case, the number of sugars contained
in the complex carbohydrate having the sugars may be 10 or
below, or 6 or below.

As specific processes for producing the complex
carbohydrates,

(1) lactose can be formed from orotic acid, galactose and
glucose by carrying out an enzyme reaction using culture
broths of a microorganism having the ability to express
Neisseria-derived β 1,4-galactosyltransferase (WO 96/10086), a
microorganism having the ability to produce UTP from a
precursor of UTP and a microorganism having the ability to
produce UDP-Gal from a sugar and UTP, or treated products of
these culture broths, as enzyme sources,

(2) N-acetyllactosamine can be formed from orotic acid,
galactose and N-acetylglucosamine by carrying out an enzyme
reaction using culture broths of a microorganism having the
ability to express Neisseria-derived

01,4-galactosyltransferase (WO 96/10086), a microorganism having the ability to produce UTP from a precursor of UTP and a microorganism having the ability to produce UDP-Gal from a sugar and UTP, or treated products of these culture broths, as enzyme sources,

(3) 3'-sialyllactose can be formed from orotic acid, N-acetylmannosamine, pyruvic acid and lactose by carrying out an enzyme reaction using culture broths of a microorganism having the ability to express *Neisseria*-derived α 2,3-sialyltransferase (*J. Biol. Chem.*, 271, 28271 (1996)), a microorganism having the ability to produce CTP from a precursor of CTP and a microorganism having the ability to produce CMP-NeuAc from a sugar and CTP, or treated products of these culture broths, as enzyme sources,

(4) 3'-sialyl-N-acetylglucosamine can be formed from orotic acid, N-acetylmannosamine, pyruvic acid and N-acetylglucosamine by carrying out an enzyme reaction using culture broths of a microorganism having the ability to express *Neisseria*-derived α 2,3-sialyltransferase (*J. Biol. Chem.*, 271, 28271 (1996)), a microorganism having the ability to produce CTP from a precursor of CTP and a microorganism having the ability to produce CMP-NeuAc from a sugar and CTP, or treated products of these culture broths, as enzyme sources,

(5) 6'-sialyl-N-acetylactosamine can be formed from orotic acid, N-acetylmannosamine, pyruvic acid and N-acetylactosamine by carrying out an enzyme reaction using culture broths of COS-7 cell line having the ability to express chicken-derived α 2,6-sialyltransferase (Eur. J. Biochem., 219, 375 (1994)), a microorganism having the ability to produce CTP from a precursor of CTP and a microorganism having the ability to produce CMP-NeuAc from a sugar and CTP, or treated products of these culture broths, as enzyme sources,

(6) GlcNAc β 1-3Gal β 1-4Glc can be formed from orotic acid, N-acetylglucosamine and lactose by carrying out an enzyme reaction using culture broths of a microorganism having the ability to express *Neisseria*-derived β 1,3-N-acetylglucosaminyltransferase (WO 96/10086), a microorganism having the ability to produce UTP from a precursor of UTP and a microorganism having the ability to produce UDP-GlcNAc from a sugar and UTP, or treated products of these culture broths, as enzyme sources,

(7) lacto-N-tetraose can be formed from orotic acid, galactose and GlcNAc β 1-3Gal β 1-4Glc by carrying out an enzyme reaction using culture broths of human melanoma cell line WM266-4 having the ability to produce β 1,3-galactosyltransferase (ATCC CRL 1676) or a transformant such as of namalwa cell line KJM-1 having the ability to

express β 1,3-galactosyltransferase gene derived from human melanoma cell line WM266-4 (Japanese Published Unexamined Patent Application No. 181759/94), a microorganism having the ability to produce UTP from a precursor of UTP and a microorganism having the ability to produce UDP-Gal from a sugar and UTP, or treated products of these culture broths, as enzyme sources,

(8) lacto-N-neotetraose can be formed from orotic acid, galactose and GlcNAc β 1-3Gal β 1-4Glc by carrying out an enzyme reaction using culture broths of *Escherichia coli* (EMBO J., 2, 3171 (1990)) or *Saccharomyces cerevisiae* (Biochem. Biophys. Res. Commun., 201, 160 (1994)) having the ability to express β 1,4-galactosyltransferase gene derived from human HeLa cell line, a microorganism having the ability to produce UTP from a precursor of UTP and a microorganism having the ability to produce UDP-Gal from a sugar and UTP, or treated products of these culture broths, as enzyme sources,

(9) lacto-N-neotetraose can be formed from orotic acid, galactose and GlcNAc β 1-3Gal β 1-4Glc by carrying out an enzyme reaction using culture broths of a microorganism having the ability to express *Neisseria*-derived β 1,4-galactosyltransferase (WO 96/10086), a microorganism having the ability to produce UTP from a precursor of UTP and a microorganism having the ability to produce UDP-Gal from a

(12) lacto-N-fucopentaose III can be formed from GMP, mannose and lacto-N-neotetraose by carrying out an enzyme reaction using culture broths of namalwa cell line having the ability to express human-derived α 1,3-fucosyltransferase (*J. Biol. Chem.*, 269, 14730 (1994)), a microorganism having the ability to produce GTP from a precursor of GTP and a microorganism having the ability to produce GDP-Fuc from a sugar and GTP, or treated products of these culture broths, as enzyme sources,

(13) lacto-N-fucopentaose III can be formed from GMP, mannose and lacto-N-neotetraose by carrying out an enzyme reaction using culture broths of a microorganism having the ability to express *Helicobacter pylori*-derived α 1,3-fucosyltransferase (*J. Biol. Chem.*, 272, 21349 and 21357 (1997)), a microorganism having the ability to produce GTP from a precursor of GTP and a microorganism having the ability to produce GDP-Fuc from a sugar and GTP, or treated products of these culture broths, as enzyme sources,

(14) globotriose can be formed from orotic acid, galactose and lactose by carrying out an enzyme reaction using culture broths of a microorganism having the ability to express *Neisseria*-derived α 1,4-galactosyltransferase (WO 96/10086), a microorganism having the ability to produce UTP from a precursor of UTP and a microorganism having the ability to

produce UDP-Gal from a sugar and UTP, or treated products of these culture broths, as enzyme sources,

(15) globotriose can be formed from orotic acid, galactose and glucose by carrying out an enzyme reaction using culture broths of a microorganism having the ability to express *Neisseria*-derived β 1,4-galactosyltransferase (WO 96/10086), a microorganism having the ability to express *Neisseria*-derived α 1,4-galactosyltransferase (WO 96/10086), a microorganism having the ability to produce UTP from a precursor of UTP and a microorganism having the ability to produce UDP-Gal from a sugar and UTP, or treated products of these culture broths, as enzyme sources,

(16) Gal α 1-4Gal β 1-4GlcNAc can be formed from orotic acid, galactose and N-acetylglucosamine by carrying out an enzyme reaction using culture broths of a microorganism having the ability to express *Neisseria*-derived α 1,4-galactosyltransferase (WO 96/10086), a microorganism having the ability to produce UTP from a precursor of UTP and a microorganism having the ability to produce UDP-Gal from a sugar and UTP, or treated products of these culture broths, as enzyme sources,

(17) lacto-N-biose can be formed from orotic acid, galactose and N-acetylglucosamine by carrying out an enzyme reaction using culture broths of an animal cell line having the ability to express human-derived

01,3-galactosyltransferase (Japanese Published Unexamined Patent Application No. 181759/94), a microorganism having the ability to produce UTP from a precursor of UTP and a microorganism having the ability to produce UDP-Gal from a sugar and UTP, or treated products of these culture broths, as enzyme sources,

02
03
04
05
06
07
08
09
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215
2216
2217
2218
2219
2220
2221
2222
2223
2224

(20) sialyl-Lewis a can be formed from GMP, mannose and sialyllacto-N-biose by carrying out an enzyme reaction using human $\alpha 1,3/1,4$ -fucosyltransferase (Carbohydrate Research, 190, 1 (1989)) and culture broths of a microorganism having the ability to produce GTP from a precursor of GTP and a microorganism having the ability to produce GDP-Fuc from a sugar and GTP, or treated products of these culture broths, as enzyme sources, and

(21) Man $\alpha 1$ -2Man can be formed from GMP and mannose by carrying out an enzyme reaction using culture broths of *Escherichia coli* having the ability to express yeast-derived $\alpha 1,2$ -mannosyltransferase (J. Org. Chem., 58, 3985 (1993)), a microorganism having the ability to produce GTP from a precursor of GTP and a microorganism having the ability to produce GDP-Man from a sugar and GTP, or treated products of these culture broths, as enzyme sources.

Processes for producing complex carbohydrates are not limited to the above-mentioned examples, and any other sugar chain can be produced industrially using a nucleotide precursor, a sugar and a complex carbohydrate precursor as the sole starting materials, within the range of glycosyltransferases which can be combined with the sugar nucleotide production process described herein and of the substrate specificity acceptable by the enzymes.

Examples of the complex carbohydrate to be produced by the production process of the present invention include

- (1) complex carbohydrates involved in the infection with pathogenic microorganisms and viruses, such as complex carbohydrates which are recognized as receptors of pathogenic microorganisms and viruses,
- (2) complex carbohydrates which are recognized as receptors of toxins produced by pathogenic microorganisms and viruses,
- (3) complex carbohydrates which are concerned, for example, in cell adhesion, recognition of foreign substances and binding of various types of lymphokine in the living body, and complex carbohydrates which contain one or a plurality of sugars such as glucose, galactose, N-acetylglucosamine, N-acetylgalactosamine, glucuronic acid, mannose, N-acetylmannosamine, fucose, sialic acid and the like, in a chemically acceptable binding mode. More specific examples include
 - (1) complex carbohydrates which are contained in the milk of human and animals, and involved in the protection of infants from microbial infection, for example, complex carbohydrates, such as lacto-N-tetraose, lacto-N-neotetraose, and the like,
 - (2) receptor complex carbohydrates which recognize microorganisms, such as *Escherichia coli*, *Propionibacterium*

granulosum, *Mycobacterium tuberculosis*, *Moraxella catarrhalis*, *Candida albicans*, *Staphylococcus saprophyticus*, *Streptococcus pneumoniae*, *Streptococcus agalactiae*, *Pseudomonas aeruginosa*, *Actinomyces naeslundii*, *Neisseria gonorrhoeae*, *Helicobacter pylori*, *Haemophilus influenzae*, and the like,

(3) receptor complex carbohydrates of viruses, such as influenza virus, coronavirus, Sendai virus, newcastle disease virus, reovirus, rotavirus, AIDS (HIV) virus, and the like,

(4) receptor complex carbohydrates of protozoa, such as *Cryptosporidium*, *Trypanosoma*, and the like,

(5) receptor complex carbohydrates having the affinity for toxins, such as cholera toxin, *Escherichia coli* heat-labile toxin, botulinum toxin, clostridial δ toxin, clostridial A toxin, Shiga toxin, Vero toxin, Shiga toxin-like toxin, *Vibrio parahaemolyticus* heat-stable toxin, tetanus toxin, and the like,

(6) cancer-related complex carbohydrates such as gangliosides (for example, GD3, GM3, etc.), globoside glycolipids, and the like,

(7) complex carbohydrates which are concerned in the adhesion of leukocytes to inflammatory regions and modification of their functions, such as sialyl-Lewis X sugar chain, and the like,

(8) complex carbohydrates concerned in autoimmune diseases, such as rheumatoid arthritis, IgA glomerulonephritis, and the like, and

(9) complex carbohydrates which are recognized by various lectin-like substances concerned in the recognition of foreign bodies and cancer cells.

Determination of the complex carbohydrate formed in the aqueous medium can be carried out in accordance with known methods (Proc. Natl. Acad. Sci. USA., 85, 3289 (1988), Anal. Biochem., 174, 459 (1988)).

Recovery of the complex carbohydrate formed in the reaction solution can be carried out in the usual way using activated carbon, an ion exchange resin and the like, for example, N-acetylglucosamine can be recovered in accordance with the process described in J. Org. Chem., 47, 5416 (1982).

Examples of the present invention are given below by way of illustration and not by way of limitation.

BEST MODE OF CARRYING OUT THE INVENTION

Example 1: Construction of recombinant plasmid capable of expressing galU and ppa

Construction process of recombinant plasmid pNT12 capable of expressing galU and ppa is described in the following (Figs. 1 and 2).

1) Construction of expression vector containing P_L promoter

Construction of pPA31 and pPAC31 as P_L promoter-containing expression vectors were carried out in the following manner (Fig. 1).

Escherichia coli JM109/pTrS30 (FERM BP-5407) which has tryptophan promoter-containing plasmid pTrS30 and another *Escherichia coli* which has P_L promoter-containing plasmid pPA1 (Japanese Published Unexamined Patent Application No. 233798/88) and P_L promoter- and cI857 repressor-containing plasmid pPAC1 (FERM BP-6054) were separately inoculated into LB medium (10 g/l Bacto-Tryptone (manufactured by Difco), 5 g/l Yeast Extract (manufactured by Difco) and 5 g/l NaCl, adjusted to pH 7.2) and cultured at 30°C for 18 hours.

From the cells obtained by the culturing, pTrS30, pPA1 and pPAC1 plasmid DNAs were isolated and purified by the above-mentioned known processes.

A 0.2 μ g portion of the thus purified pTrS30 DNA was cleaved with restriction enzymes *Pst*I and *Cla*I, the resulting DNA fragments were separated by agarose gel electrophoresis and then a fragment of 3.4 kb was recovered using Gene Clean II Kit (manufactured by Bio101). A 0.5 μ g portion of the purified pPA1 DNA was cleaved with restriction enzymes *Pst*I and *Cla*I, the resulting DNA fragments were separated by agarose gel electrophoresis and then a fragment of 1.0 kb was recovered in the same manner.

Using a ligation kit (TAKARA Ligation Kit, manufactured by Takara Shuzo Co., Ltd.), the fragments of 3.4 kb and 1.0 kb were subjected to ligation reaction for 16 hours at 16°C.

Using the ligation reaction solution, *Escherichia coli* NM522 was transformed in accordance with the above-mentioned known process, and the resulting transformant was spread on LB agar medium containing 50 µg/ml ampicillin and then cultured overnight at 37°C.

By extracting a plasmid from the thus grown colonies of the transformant in accordance with the above-mentioned known process, the P_L promoter-effected expression vector pPA31 was obtained. Structure of the plasmid was recognized by restriction enzyme cleavage (Fig. 1).

A 0.2 µg portion of the purified pPA31 DNA was cleaved with restriction enzymes *Pst*I and *Cla*I, the resulting DNA fragments were separated by agarose gel electrophoresis and then a fragment of 3.4 kb was recovered using Gene Clean II Kit. A 0.5 µg portion of the purified pFAC1 DNA was cleaved with restriction enzymes *Pst*I and *Cla*I, the resulting DNA fragments were separated by agarose gel electrophoresis and then a fragment of 2.3 kb was recovered in the same manner.

Using a ligation kit, the fragments of 3.4 kb and 2.3 kb were subjected to ligation reaction for 16 hours at 16°C.

Using the ligation reaction solution, *Escherichia coli* NM522 was transformed in accordance with the above-mentioned known process, and the resulting transformant was spread on LB agar medium containing 50 µg/ml ampicillin and then cultured overnight at 37°C.

By extracting a plasmid from the thus grown colonies of the transformant in accordance with the above-mentioned known process, the P_i promoter-effected cI857 repressor-containing expression vector pPAC31 was obtained. Structure of the plasmid was recognized by restriction enzyme cleavage (Fig. 1).

2) Construction of galU expression plasmid

Chromosomal DNA of *Escherichia coli* W3100 was isolated and purified by a known process (for example, Current Protocols in Molecular Biology, John Wiley and Sons Inc. (1994)).

The sense strand DNA primer shown in SEQ ID NO:1 and the antisense strand DNA primer shown in SEQ ID NO:2 were synthesized using 380A DNA Synthesizer manufactured by Applied Biosystems.

The PCR process was carried out using the synthesized DNA strands as primers, and the chromosomal DNA of the strain W3110 as the template. The PCR was effected using 40 µl of a reaction solution containing 0.04 µg of the W3110 chromosomal DNA, 0.5 µM of each primer, 1.0 unit of TAKARA Ex Taq

(manufactured by Takara Shuzo Co., Ltd.), 4 μ l of 10 \times Ex Taq buffer (manufactured by Takara Shuzo Co., Ltd.) and 200 μ M of each deoxyNTP, and repeating 30 cycles of the reaction, each cycle containing 94°C for 1 minute, 42°C for 2 minutes and 72°C for 3 minutes.

A 1/10 volume of the reaction solution was subjected to agarose gel electrophoresis to verify amplification of the fragment of interest, and then the remaining reaction solution was mixed with the same volume of phenol/chloroform (1 vol/1 vol) solution with saturated TE (10 mM Tris-HCl buffer (pH 8.0) and 1 mM EDTA). The mixture solution was centrifuged, and the thus obtained upper layer was mixed with 2 volumes of cold ethanol and allowed to stand for 30 minutes at -80°C. The solution after standing was centrifuged to obtain a precipitate of DNA. The precipitate was washed with 70% cold ethanol and dried in vacuo to recover the precipitate. Hereinafter, the steps starting from the addition of phenol/chloroform solution with saturated TE until the recovery of the ethanol-washed DNA are referred to as an ethanol precipitation process.

The DNA precipitate was dissolved in 20 μ l of TE. Using 5 μ l portion of the solution, DNA was cleaved with restriction enzymes *Hind*III and *Bam*HI, the resulting DNA fragments were separated by agarose gel electrophoresis and then a fragment of 0.9 kb was recovered using Gene Clean II

Kit. A 0.2 µg portion of pPA31 DNA obtained in Example 1-1) was cleaved with restriction enzymes *HindIII* and *BamHI*, the resulting DNA fragments were separated by agarose gel electrophoresis and then a fragment of 4.2 kb was recovered in the same manner.

Using a ligation kit, the fragments of 0.9 kb and 4.2 kb were subjected to ligation reaction for 16 hours at 16°C.

Using the ligation reaction solution, *Escherichia coli* KY8415 was transformed in accordance with the above-mentioned known process, and the resulting transformant was spread on LB agar medium containing 50 µg/ml ampicillin and then cultured overnight at 30°C.

By extracting a plasmid from the thus grown colonies of the transformant in accordance with the above-mentioned known process, galU expression plasmid pNT9 was obtained. Structure of the plasmid was recognized by restriction enzyme cleavage (Fig. 2).

3) Construction of simultaneous galU, ppa expression plasmid

The sense strand DNA primer shown in SEQ ID NO:3 and the antisense strand DNA primer shown in SEQ ID NO:4 were synthesized, and the PCR was carried out using the synthesized DNA strands as primers, and the chromosomal DNA of the strain W3110 as the template, under the same conditions as described in the foregoing.

After completion of the PCR treatment, a precipitate of DNA was obtained by the ethanol precipitation process. The precipitate was dissolved in 20 μ l of TE. Using 5 μ l portion of the solution, the DNA was cleaved with restriction enzymes *Bam*HI and *Sal*I, the resulting DNA fragments were separated by agarose gel electrophoresis and then a fragment of 1.0 kb was recovered using Gene Clean II Kit. A 0.2 μ g portion of pNT9 DNA obtained in Example 1-2) was cleaved with restriction enzymes *Bam*HI and *Sal*I, the resulting DNA fragments were separated by agarose gel electrophoresis and then a fragment of 4.9 kb was recovered in the same manner.

Using a ligation kit, the fragments of 1.0 kb and 4.9 kb were subjected to ligation reaction for 16 hours at 16°C.

Using the ligation reaction solution, *Escherichia coli* KY8415 was transformed in accordance with the above-mentioned known process, and the resulting transformant was spread on LB agar medium containing 50 μ g/ml ampicillin and then cultured overnight at 30°C.

By extracting a plasmid from the thus grown colonies of the transformant in accordance with the above-mentioned known process, the *gal*U, *ppa* simultaneous expression plasmid pNT12 was obtained. Structure of the plasmid was recognized by restriction enzyme cleavage (Fig. 2).

A 0.5 μ g portion of the pNT12 DNA was cleaved with restriction enzymes *Eco*RI and *Sal*I, the resulting DNA

fragments were separated by agarose gel electrophoresis and then a fragment of 2.2 kb was recovered using Gene Clean II Kit. Separately from this, a 0.2 µg portion of pSTV28 DNA (manufactured by Takara Shuzo Co., Ltd.) was cleaved with restriction enzymes *EcoRI* and *SalI*, the resulting DNA fragments were separated by agarose gel electrophoresis and then a fragment of 3.0 kb was recovered in the same manner.

Using a ligation kit, the fragments of 2.2 kb and 3.0 kb were subjected to ligation reaction for 16 hours at 16°C.

Using the ligation reaction solution, *Escherichia coli* NM522 was transformed in the usual way, and the resulting transformant was spread on LB agar medium containing 10 µg/ml chloramphenicol and then cultured overnight at 30°C.

By extracting a plasmid from the thus grown colonies of the transformant in the usual way, the galU, ppa simultaneous expression plasmid pNT32 was obtained. Structure of the plasmid was recognized by restriction enzyme cleavage (Fig. 2).

Example 2: Production of UDP-Glc

Escherichia coli KY8415/pNT12 obtained in Example 1 was inoculated into a 1 L baffled conical flask containing 125 ml of LB medium supplemented with 50 µg/ml ampicillin and cultured at 30°C for 17 hours under shaking at 220 rpm. A

125 ml portion of the culture broth was inoculated into a 5 L culture vessel containing 2.5 L of an aqueous medium (without adjusting pH) which contains 10 g/l glucose, 12 g/l Bacto-Tryptone (manufactured by Difco), 24 g/l Yeast Extract (manufactured by Difco), 2.3 g/l KH_2PO_4 (separate sterilization), 12.5 g/l K_2HPO_4 (separate sterilization) and 50 $\mu\text{g/ml}$ ampicillin and cultured at 30°C for 4 hours and then at 40°C for 3 hours, under conditions of 600 rpm and 2.5 L/min of aeration.

During the culturing, pH of the medium was maintained at 7.0 using 28% aqueous ammonia. In addition, glucose was added during the culturing when required. The culture broth was centrifuged to obtain wet cells. As occasion demands, these wet cells can be preserved at -20°C and utilized by thawing the cells prior to use.

Corynebacterium ammoniagenes ATCC 21170 was inoculated into a 300 ml-baffled conical flask containing 20 ml of an aqueous medium of 50 g/l glucose, 10 g/l Polypeptone (manufactured by Nippon Seiyaku), 10 g/l Yeast Extract (manufactured by Oriental Yeast), 5 g/l urea, 5 g/l $(\text{NH}_4)_2\text{SO}_4$, 1 g/l KH_2PO_4 , 3 g/l K_2HPO_4 , 1 g/l $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.1 g/l $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$, 10 mg/l $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 10 mg/l $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$, 20 mg/l $\text{MnSO}_4 \cdot 4\text{H}_2\text{O}$, 20 mg/l L-cysteine, 10 mg/l calcium D-pantothenate, 5 mg/l vitamin B_1 , 5 mg/l nicotinic acid and 30 $\mu\text{g/l}$ biotin

(adjusted to pH 7.2 with 10 N NaOH) and cultured at 28°C for 24 hours under shaking at 220 rpm.

A 20 ml portion of the culture broth was inoculated into a 2 L baffled conical flask containing 250 ml of the same aqueous medium and cultured at 28°C for 24 hours under shaking at 220 rpm. The thus obtained culture broth was used as a seed culture broth.

A 250 ml portion of the seed culture broth was inoculated into a 5 L culture vessel containing 2.25 L of an aqueous medium of 150 g/l glucose, 5 g/l meat extract (manufactured by Kyokuto Pharmaceutical Industry), 10 g/l KH_2PO_4 , 10 g/l K_2HPO_4 , 10 g/l $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.1 g/l $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$, 20 mg/l $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 10 mg/l $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$, 20 mg/l $\text{MnSO}_4 \cdot 4\text{H}_2\text{O}$ (separate sterilization), 15 mg/l β -alanine (separate sterilization), 20 mg/l L-cysteine, 100 $\mu\text{g/l}$ biotin, 2 g/l urea and 5 mg/l vitamin B_1 (separate sterilization) (adjusted to pH 7.2 with 10 N NaOH) and cultured at 32°C for 24 hours under conditions of 600 rpm and 2.5 L/min of aeration. During the culturing, pH of the culture broth was maintained at 6.8 using 28% aqueous ammonia.

The culture broth was centrifuged to obtain wet cells. As occasion demands, these wet cells can be preserved at -20°C and utilized by thawing the cells prior to use.

A 30 ml portion of a reaction solution having a composition of 40 g/l *Escherichia coli* KY8415/pNT12 wet cells,

150 g/l *Corynebacterium ammoniagenes* ATCC 21170 wet cells, 100 g/l glucose, 20 g/l KH_2PO_4 , 5 g/l $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 5 g/l phytic acid, 21.2 g/l orotic acid (potassium salt), 4 g/l Nymean S-215 and 10 ml/l xylene was put into a 200 ml beaker, and 21 hours of the reaction was carried out at 32°C under stirring the reaction solution with a magnetic stirrer (900 rpm).

During the reaction, pH of the reaction solution was maintained at 7.2 using 4 N NaOH, and glucose and KH_2PO_4 were added when required.

By the reaction, 43.9 g/l UDP-Glc (2Na salt) was formed in the reaction solution.

Example 3: Construction of recombinant plasmid capable of expressing galT and galK

Construction process of recombinant plasmid pMT25 capable of expressing galT and galK is described in the following (Fig. 3).

The sense strand DNA primer shown in SEQ ID NO:5 and the antisense strand DNA primer shown in SEQ ID NO:6 were synthesized. The PCR was carried out using the synthesized DNA strands as primers, and the chromosomal DNA of the strain W3110 as the template, under the same conditions as described in the foregoing.

After completion of the PCR treatment, a precipitate of DNA was obtained by the ethanol precipitation process.

[illegible]

1. **Author(s)**
 2. **Title**
 3. **Journal**
 4. **Volume**
 5. **Issue**
 6. **Page(s)**
 7. **Year**
 8. **DOI**
 9. **URL**
 10. **Accession Number**
 11. **Keywords**
 12. **Abstract**
 13. **Notes**
 14. **References**
 15. **Comments**
 16. **History**
 17. **Metadata**
 18. **Full Text**
 19. **Download**
 20. **Print**
 21. **Share**
 22. **Feedback**
 23. **Help**
 24. **Search**
 25. **Advanced Search**
 26. **Help**
 27. **Feedback**
 28. **Help**
 29. **Feedback**
 30. **Help**
 31. **Feedback**
 32. **Help**
 33. **Feedback**
 34. **Help**
 35. **Feedback**
 36. **Help**
 37. **Feedback**
 38. **Help**
 39. **Feedback**
 40. **Help**
 41. **Feedback**
 42. **Help**
 43. **Feedback**
 44. **Help**
 45. **Feedback**
 46. **Help**
 47. **Feedback**
 48. **Help**
 49. **Feedback**
 50. **Help**
 51. **Feedback**
 52. **Help**
 53. **Feedback**
 54. **Help**
 55. **Feedback**
 56. **Help**
 57. **Feedback**
 58. **Help**
 59. **Feedback**
 60. **Help**
 61. **Feedback**
 62. **Help**
 63. **Feedback**
 64. **Help**
 65. **Feedback**
 66. **Help**
 67. **Feedback**
 68. **Help**
 69. **Feedback**
 70. **Help**
 71. **Feedback**
 72. **Help**
 73. **Feedback**
 74. **Help**
 75. **Feedback**
 76. **Help**
 77. **Feedback**
 78. **Help**
 79. **Feedback**
 80. **Help**
 81. **Feedback**
 82. **Help**
 83. **Feedback**
 84. **Help**
 85. **Feedback**
 86. **Help**
 87. **Feedback**
 88. **Help**
 89. **Feedback**
 90. **Help**
 91. **Feedback**
 92. **Help**
 93. **Feedback**
 94. **Help**
 95. **Feedback**
 96. **Help**
 97. **Feedback**
 98. **Help**
 99. **Feedback**
 100. **Help**

[illegible]

A 0.2 µg portion of the pPAC31 DNA obtained in Example 1-1) was cleaved with restriction enzymes *Cla*I and *Bam*HI, the resulting DNA fragments were separated by agarose gel electrophoresis and then a fragment of 5.5 kb was recovered in the same manner.

Using a ligation kit, the fragments of 2.3 kb and 5.5 kb were subjected to ligation reaction for 16 hours at 16°C.

Using the ligation reaction solution, *Escherichia coli* NM52 was transformed in accordance with the above-mentioned known process, and the resulting transformant was spread on LB agar medium containing 50 µg/ml ampicillin and then cultured overnight at 30°C.

By extracting a plasmid from the thus grown colonies of the transformant in accordance with the above-mentioned known process, plasmid pNT25 capable of expressing *galT* and *galK* simultaneously was obtained. Structure of the plasmid was recognized by restriction enzyme cleavage (Fig. 3).

Example 4: Production of UDP-Gal

1) Preparation of *galT*, *galK*, *galU*, *ppa* expression strain

Using the pNT32 DNA obtained in Example 1-3), *Escherichia coli* NM522/pNT25 was transformed in accordance with the above-mentioned known process, and the resulting transformant was spread on LB agar medium containing 50 µg/ml ampicillin and 10 µg/ml chloramphenicol and then cultured

overnight at 30°C. By selecting the thus grown transformants, *Escherichia coli* NM522/pNT25/pNT32 was obtained as the galT, galK, galU, ppa expression strain.

2) Production of UDP-Gal

Escherichia coli NM522/pNT25/pNT32 obtained in Example 4-1) was cultured in the same manner as in Example 2, and the culture broth was centrifuged to obtain wet cells. Also, *Corynebacterium ammoniagenes* ATCC 21170 was cultured in the same manner as in Example 2, and the culture broth was centrifuged to obtain wet cells. As occasion demands, the wet cells can be preserved at -20°C and utilized by thawing them prior to use.

A 2 L portion of a reaction solution having a composition of 50 g/l *Escherichia coli* NM522/pNT25/pNT32 wet cells, 150 g/l *Corynebacterium ammoniagenes* ATCC 21170 wet cells, 80 g/l glucose, 20 g/l galactose, 15 g/l KH_2PO_4 , 5 g/l $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 5 g/l phytic acid, 21.2 g/l crotonic acid (potassium salt), 4 g/l Nymean S-215 and 10 ml/l xylene was put into 5 L culture vessel, and 26 hours of the reaction was carried out at 32°C under stirring the reaction solution at 600 rpm with an aeration rate of 1 L/min.

During the reaction, pH of the reaction solution was maintained at 7.2 using 4 N NaOH, and glucose, galactose and KH_2PO_4 were added when required.

By the reaction, 47.4 g/l UDP-Gal (2Na salt) was formed in the reaction solution.

Example 5: Construction of recombinant plasmid capable of expressing galT and galK in *Corynebacterium ammoniagenes*

Construction process of recombinant plasmid pTK7 capable of expressing *Escherichia coli*-derived galT and galK in *Corynebacterium ammoniagenes* is described in the following (Fig. 4).

1) Construction of pCG116

Plasmid pCG116 capable of replicating in *Corynebacterium ammoniagenes* was constructed in the following manner.

A 0.5 µg portion of plasmid pCG11 (Japanese Published Examined Patent Application No. 91827/94) DNA was cleaved with restriction enzymes *Pst*I and *Stu*I, the resulting DNA fragments were separated by agarose gel electrophoresis and then a fragment of 6.5 kb was recovered using Gene Clean II Kit.

On the other hand, a 1.0 µg portion of plasmid pUC19 DNA was cleaved with a restriction enzyme *Eco*RI and then blunt-ended using DNA Blunting Kit (manufactured by Takara Shuzo Co., Ltd.). The DNA thus blunt-ended was cleaved with a restriction enzyme *Pst*I, the resulting DNA fragments were separated by agarose gel electrophoresis and then a fragment

of 43 bp was recovered using MERmaid Kit (manufactured by Biol01).

Using a ligation kit, the fragments of 6.5 kb and 43 bp were subjected to ligation reaction for 16 hours at 16°C.

Using the ligation reaction solution, *Corynebacterium ammoniagenes* ATCC 21170 was transformed by the electroporation method (FEMS Microbiol. Lett., 65, 299 (1989)), and the resulting transformant was spread on LB agar medium containing 100 µg/ml spectinomycin and then cultured at 30°C for 2 days.

By extracting a plasmid from the thus grown colonies of the transformant in accordance with the known process (J. Bacteriol., 159, 306 (1984)), plasmid pCG116 was obtained. Structure of the plasmid was recognized by restriction enzyme cleavage (Fig. 4).

2) Construction of pTK7 capable of expressing galT and galK

A 1.0 µg portion of the galT and galK expression plasmid pNT25 DNA obtained in Example 3 was cleaved with restriction enzymes XhoI and BamHI, the resulting DNA fragments were separated by agarose gel electrophoresis and then a fragment of 3.5 kb was recovered using Gene Clean II Kit.

On the other hand, a 0.5 µg portion of the plasmid pCG116 DNA prepared in Example 5-1) was cleaved with a restriction enzymes SalI and BamHI, the resulting DNA

fragments were separated by agarose gel electrophoresis and then a fragment of 6.5 kb was recovered in the same manner.

Using a ligation kit, the fragments of 3.5 kb and 6.5 kb were subjected to ligation reaction for 16 hours at 16°C.

Using the ligation reaction solution, *Corynebacterium ammoniagenes* ATCC 21170 was transformed by the electroporation process, and the resulting transformant was spread on LB agar medium containing 100 µg/ml spectinomycin and then cultured at 30°C for 2 days.

By extracting a plasmid from the thus grown colonies of the transformant in accordance with the known process, plasmid pTK7 capable of expressing galT and galK simultaneously was obtained. Structure of the plasmid was recognized by restriction enzyme cleavage (Fig. 4).

Example 6: Production of UDP-Gal

Corynebacterium ammoniagenes ATCC 21170/pTK7 obtained in Example 5 was cultured by the same process as described in Example 2 at 32°C for 20 hours and then at 40°C for 4 hours, and the thus obtained culture broth was centrifuged to obtain wet cells. As occasion demands, the wet cells can be preserved at -20°C and utilized by thawing them prior to use.

A 30 ml portion of a reaction solution having a composition of 150 g/l *Corynebacterium ammoniagenes* ATCC 21170/pTK7 wet cells, 40 g/l fructose, 20 g/l galactose, 15

g/l KH_2PO_4 , 5 g/l $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 5 g/l phytic acid, 10.6 g/l orotic acid (potassium salt), 4 g/l Nymeen S-215 and 10 ml/l xylene was put into a 200 ml beaker, and 22 hours of the reaction was carried out at 32°C under stirring the reaction solution with a magnetic stirrer (900 rpm).

During the reaction, pH of the reaction solution was maintained at 7.2 using 4 N NaOH, and fructose, galactose and KH_2PO_4 were added when required.

By the reaction, 7.2 g/l UDP-Gal (2Na salt) was formed in the reaction solution.

Example 7: Construction of glmU, ppa, pgm, glmM, glk and pfkB expression plasmid

1) Construction of glmU and ppa expression plasmid

The sense strand DNA primer shown in SEQ ID NO:7 and the antisense strand DNA primer shown in SEQ ID NO:8 were synthesized. The PCR was carried out using the synthesized DNA strands as primers, and the chromosomal DNA of the strain W3110 as the template, under the same conditions as described in the foregoing.

After completion of the PCR treatment, a precipitate of DNA was obtained by the ethanol precipitation process. The DNA precipitate was dissolved in 20 μl of TE. Using 5 μl portion of the solution, the DNA was cleaved with restriction enzymes HindIII and BamHI, the resulting DNA fragments were

separated by agarose gel electrophoresis and then a fragment of 1.4 kb was recovered using Gene Clean II Kit. A 0.5 µg portion of pFA31 DNA obtained in Example 1-1) was cleaved with restriction enzymes *HindIII* and *BamHI*, the resulting DNA fragments were separated by agarose gel electrophoresis and then a fragment of 4.2 kb was recovered in the same manner.

Using a ligation kit, the fragments of 1.4 kb and 4.2 kb were subjected to ligation reaction for 16 hours at 16°C.

Using the ligation reaction solution, *Escherichia coli* KY8415 was transformed in accordance with the above-mentioned known process, and the resulting transformant was spread on LB agar medium containing 50 µg/ml ampicillin and then cultured overnight at 30°C.

By extracting a plasmid from the thus grown colonies of the transformant in accordance with the above-mentioned known process, *glmU* expression plasmid pNT10 was obtained. Structure of the plasmid was recognized by restriction enzyme cleavage (Fig. 5).

A 0.5 µg portion of the pNT12 DNA obtained in Example 1-3) was cleaved with restriction enzymes *BamHI* and *SalI*, the resulting DNA fragments were separated by agarose gel electrophoresis and then a fragment of 1.0 kb was recovered in the same manner. A 0.2 µg portion of the just described pNT10 DNA was cleaved with restriction enzymes *BamHI* and *SalI*, the resulting DNA fragments were separated by agarose gel

electrophoresis and then a fragment of 5.3 kb was recovered in the same manner.

Using a ligation kit, the fragments of 1.0 kb and 5.3 kb were subjected to ligation reaction for 16 hours at 16°C.

Using the ligation reaction solution, *Escherichia coli* KY8415 was transformed in accordance with the above-mentioned known process, and the resulting transformant was spread on LB agar medium containing 50 µg/ml ampicillin and then cultured overnight at 30°C.

By extracting a plasmid from the thus grown colonies of the transformant in accordance with the above-mentioned known process, the glmU, ppa simultaneous expression plasmid pNT14 was obtained. Structure of the plasmid was recognized by restriction enzyme cleavage (Fig. 5).

2) Construction of pgm expression plasmid

The sense strand DNA primer shown in SEQ ID NO:9 and the antisense strand DNA primer shown in SEQ ID NO:10 were synthesized. The PCR was carried out using the synthesized DNA strands as primers, and the chromosomal DNA of the strain W3110 as the template, under the same conditions as described in the foregoing.

After completion of the PCR, a precipitate of DNA was obtained by the ethanol precipitation process. The precipitate was dissolved in 20 µl of TE. Using 5 µl portion of the solution, the DNA was cleaved with restriction enzymes

ClaI and BamHI, the resulting DNA fragments were separated by agarose gel electrophoresis and then a fragment of 1.8 kb was recovered using Gene Clean II Kit. A 0.2 µg portion of the pPAC31 DNA obtained in Example 1-1) was cleaved with restriction enzymes ClaI and BamHI, the resulting DNA fragments were separated by agarose gel electrophoresis and then a fragment of 5.5 kb was recovered in the same manner.

Using a ligation kit, the fragments of 1.8 kb and 5.5 kb were subjected to ligation reaction for 16 hours at 16°C.

Using the ligation reaction solution, *Escherichia coli* NM522 was transformed in accordance with the above-mentioned known process, and the resulting transformant was spread on LB agar medium containing 50 µg/ml ampicillin and then cultured overnight at 30°C.

By extracting a plasmid from the thus grown colonies of the transformant in accordance with the above-mentioned known process, pgm expression plasmid pNT24 was obtained. Structure of the plasmid was recognized by restriction enzyme cleavage (Fig. 6).

3) Construction of glmM expression plasmid

The sense strand DNA primer shown in SEQ ID NO:11 and the antisense strand DNA primer shown in SEQ ID NO:12 were synthesized. The PCR was carried out using the synthesized DNA strands as primers, and the chromosomal DNA of

Escherichia coli W3110 as the template, under the same conditions as described in the foregoing.

After completion of the PCR, a precipitate of DNA was obtained by the ethanol precipitation process. The precipitate was dissolved in 20 μ l of TE. Using 5 μ l portion of the solution, the DNA was cleaved with restriction enzymes *Cla*I and *Bam*HI, the resulting DNA fragments were separated by agarose gel electrophoresis and then a fragment of 1.6 kb was recovered using Gene Clean II Kit. A 0.2 μ g portion of the pPAC31 DNA obtained in Example 1-1) was cleaved with restriction enzymes *Cla*I and *Bam*HI, the resulting DNA fragments were separated by agarose gel electrophoresis and then a fragment of 5.5 kb was recovered in the same manner.

Using a ligation kit, the fragments of 1.6 kb and 5.5 kb were subjected to ligation reaction for 16 hours at 16°C.

Using the ligation reaction solution, *Escherichia coli* NM522 was transformed in accordance with the above-mentioned known process, and the resulting transformant was spread on LB agar medium containing 50 μ g/ml ampicillin and then cultured overnight at 30°C.

By extracting a plasmid from the thus grown colonies of the transformant in accordance with the above-mentioned known process, glmM expression plasmid pNT44 was obtained. Structure of the plasmid was recognized by restriction enzyme cleavage (Fig. 7).

4) Construction of glk expression plasmid

The sense strand DNA primer shown in SEQ ID NO:13 and the antisense strand DNA primer shown in SEQ ID NO:14 were synthesized. The PCR was carried out using the synthesized DNA strands as primers, and the chromosomal DNA of *Escherichia coli* W3110 as the template, under the same conditions as described in the foregoing.

After completion of the PCR, a precipitate of DNA was obtained by the ethanol precipitation process. The precipitate was dissolved in 20 μ l of TE. Using 5 μ l portion of the solution, the DNA was cleaved with restriction enzymes *Hind*III and *Bam*HI, the resulting DNA fragments were separated by agarose gel electrophoresis and then a fragment of 0.5 kb was recovered using Gene Clean II Kit.

A 0.2 μ g portion of the pPA31 DNA obtained in Example 1-1) was cleaved with restriction enzymes *Hind*III and *Bam*HI, the resulting DNA fragments were separated by agarose gel electrophoresis and then a fragment of 4.2 kb was recovered in the same manner.

Using a ligation kit, the fragments of 0.5 kb and 4.2 kb were subjected to ligation reaction for 16 hours at 16°C.

Using the ligation reaction solution, *Escherichia coli* NM522 was transformed in accordance with the usual process, and the resulting transformant was spread on LB agar

medium containing 50 µg/ml ampicillin and then cultured overnight at 30°C.

By extracting a plasmid from the thus grown colonies of the transformant in the usual way, plasmid pNT45 containing a part of glk was obtained. Structure of the plasmid was recognized by restriction enzyme cleavage (Fig. 8).

The PCR was carried out under the same conditions described above, the DNA contained in 5 µl portion of the thus obtained DNA solution was cleaved with a restriction enzyme *HindIII*, the resulting DNA fragments were separated by agarose gel electrophoresis and then a fragment of 0.5 kb was recovered in the same manner. A 0.2 µg portion of the pNT45 DNA obtained by the just described process was cleaved with the restriction enzyme *HindIII* and subjected to a dephosphorylation treatment with alkaline phosphatase, the resulting DNA fragments were separated by agarose gel electrophoresis and then a fragment of 4.7 kb was recovered in the same manner.

Using a ligation kit, the fragments of 0.5 kb and 4.7 kb were subjected to ligation reaction for 16 hours at 16°C.

Using the ligation reaction solution, *Escherichia coli* NM522 was transformed in accordance with the usual way, and the resulting transformant was spread on LB agar medium

containing 50 µg/ml ampicillin and then cultured overnight at 30°C.

By extracting a plasmid from the thus grown colonies of the transformant in accordance with the usual way, glk expression plasmid pNT46 was obtained. Structure of the plasmid was recognized by restriction enzyme cleavage (Fig. 8).

5) Construction of pfkB expression plasmid

The sense strand DNA primer shown in SEQ ID NO:15 and the antisense strand DNA primer shown in SEQ ID NO:16 were synthesized. The PCR was carried out using the synthesized DNA strands as primers, and the chromosomal DNA of the strain W3110 as the template, under the same conditions as described in the foregoing.

After completion of the PCR, a precipitate of DNA was obtained by the ethanol precipitation process. The precipitate was dissolved in 20 µl of TE. Using 5 µl portion of the solution, the DNA was cleaved with restriction enzymes *HindIII* and *EcoRV*, the resulting DNA fragments were separated by agarose gel electrophoresis and then a fragment of 1.3 kb was recovered using Gene Clean II Kit. A 0.2 µg portion of pBluescript II SK+ DNA was cleaved with restriction enzymes *HindIII* and *EcoRV*, the resulting DNA fragments were separated by agarose gel electrophoresis and then a fragment of 3.0 kb was recovered in the same manner.

Using a ligation kit, the fragments of 1.3 kb and 3.0 kb were subjected to ligation reaction for 16 hours at 16°C.

Using the ligation reaction solution, *Escherichia coli* NM522 was transformed in accordance with the usual process, and the resulting transformant was spread on LB agar medium containing 50 µg/ml ampicillin and then cultured overnight at 30°C.

By extracting a plasmid from the thus grown colonies of the transformant in accordance with the usual process, plasmid pNT43 containing the pfkB gene was obtained. Structure of the plasmid was recognized by restriction enzyme cleavage (Fig. 9).

A 0.5 µg portion of the pNT43 DNA was cleaved with restriction enzymes *Cla*I and *Sac*I, the resulting DNA fragments were separated by agarose gel electrophoresis and then a fragment of 1.3 kb was recovered in the same manner.

A 0.2 µg portion of the pPAC31 DNA obtained in Example 1-1) was cleaved with restriction enzymes *Cla*I and *Sac*I, the resulting DNA fragments were separated by agarose gel electrophoresis and then a fragment of 5.7 kb was recovered in the same manner.

Using a ligation kit, the fragments of 1.3 kb and 5.7 kb were subjected to ligation reaction for 16 hours at 16°C.

Using the ligation reaction solution, *Escherichia coli* NM522 was transformed in accordance with the usual

process, and the resulting transformant was spread on LB agar medium containing 50 µg/ml ampicillin and then cultured overnight at 30°C.

By extracting a plasmid from the thus grown colonies of the transformant in accordance with the usual process, pfkB expression plasmid pNT47 was obtained. Structure of the plasmid was recognized by restriction enzyme cleavage (Fig. 9).

Example 8: Production of UDP-GlcNAc

Escherichia coli KY8415/pNT14, NM522/pNT24, NM522/pNT44 and NM522/pNT47 obtained in Example 7 were cultured in the same manner as in Example 2, and each of the thus obtained culture broths were centrifuged to obtain wet cells. As occasion demands, these wet cells can be preserved at -20°C and utilized by thawing the cells prior to use.

A 0.1 ml portion of a reaction solution having a composition of 6 g/l *Escherichia coli* NM522/pNT24 wet cells, 6 g/l NM522/pNT47 wet cells, 100 mM Tris-HCl buffer (pH 8.0), 6 mM MgCl₂·6H₂O, 10 mM glucose-6-phosphate, 2.5 mM fructose-6-phosphate, 2.5 mM ATP and 4 g/l Nymeen S-215 was put into a 1.5 ml tube, and 1 hour of the reaction was carried out at 37°C. The reaction solution was treated at 65°C for 5 minutes and shortage in cells and substances was suspended until 0.3 g/l *Escherichia coli* KY8415/pNT14 wet cells, 6 g/l

[illegible][illegible][illegible][illegible][illegible]

[illegible][illegible][illegible][illegible][illegible][illegible]

medium containing 50 µg/ml ampicillin and then cultured overnight at 30°C.

By extracting a plasmid from the thus grown colonies of the transformant in accordance with the usual process, galK expression plasmid pNT54 was obtained. Structure of the plasmid was recognized by restriction enzyme cleavage (Fig. 10).

Example 11: Production of UDP-GlcNAc

Escherichia coli NM522/pNT54 obtained in Example 10 was cultured in the same manner as in Example 2, and the culture broth was centrifuged to obtain wet cells. Also, *Corynebacterium ammoniagenes* ATCC 21170 was cultured in the same manner as in Example 2, and the culture broth was centrifuged to obtain wet cells. As occasion demands, these wet cells can be preserved at -20°C and utilized by thawing the cells prior to use.

A 30 ml portion of a reaction solution having a composition of 50 g/l *Escherichia coli* NM522/pNT54 wet cells, 150 g/l *Corynebacterium ammoniagenes* ATCC 21170 wet cells, 40 g/l fructose, 67 g/l N-acetylglucosamine, 15 g/l KH_2PO_4 , 5 g/l $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 5 g/l phytic acid, 10 g/l orotic acid (potassium salt), 4 g/l Nymeen S-215 and 10 ml/l xylene was put into a 200 ml beaker, and 27 hours of the reaction was

carried out at 32°C under stirring the reaction solution with a magnetic stirrer (900 rpm).

During the reaction, pH of the reaction solution was maintained at 7.2 using 4 N NaOH, and fructose and KH_2PO_4 were added when required.

By the reaction, 17.1 g/l UDP-GlcNAc (2Na salt) was formed in the reaction solution.

Example 12: Simultaneous production of UDP-GlcNAc and UDP-Gal

The strain NM522/pNT25 obtained in Example 3 was cultured in the same manner as in Example 2, and the culture broth was centrifuged to obtain wet cells. Also, *Corynebacterium ammoniagenes* ATCC 21170 was cultured in the same manner as in Example 2, and the culture broth was centrifuged to obtain wet cells. As occasion demands, these wet cells can be preserved at -20°C and utilized by thawing the cells prior to use.

A 30 ml portion of a reaction solution having a composition of 25 g/l *Escherichia coli* NM522/pNT25 wet cells, 150 g/l *Corynebacterium ammoniagenes* ATCC 21170 wet cells, 60 g/l fructose, 50 g/l N-acetylglucosamine, 40 g/l galactose, 15 g/l KH_2PO_4 , 5 g/l $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 5 g/l phytic acid, 10 g/l orotic acid (potassium salt), 4 g/l Nymeen S-215 and 10 ml/l xylene was put into a 200 ml beaker, and 24 hours of the

reaction was carried out at 32°C under stirring the reaction solution with a magnetic stirrer (900 rpm).

During the reaction, pH of the reaction solution was maintained at 7.2 using 4 N NaOH, and KH_2PO_4 was added when required.

By the reaction, 11.4 g/l UDP-GlcNAc (2Na salt) and 18 g/l UDP-Gal (2Na salt) were formed in the reaction solution.

Example 13: Construction of plasmid capable of expressing *manB*, *manC*, *pgm* and *pfkB*

1) Construction of *manB*, *manC* expression plasmid

The sense strand DNA primer shown in SEQ ID NO:17 and the antisense strand DNA primer shown in SEQ ID NO:18 were synthesized. The PCR was carried out using the synthesized DNA strands as primers, and the chromosomal DNA of *Escherichia coli* W3110 as the template, under the same conditions as described in the foregoing.

After completion of the PCR treatment, a precipitate of DNA was obtained by the ethanol precipitation process. The precipitate was dissolved in 20 µl of TE. Using 5 µl portion of the solution, the DNA was cleaved with restriction enzymes *HindIII* and *BamHI*, the resulting DNA fragments were separated by agarose gel electrophoresis and then a fragment of 3.0 kb was recovered using Gene Clean II Kit. A 0.2 µg

portion of pBluescript II SK+ DNA was cleaved with restriction enzymes HindIII and BamHI, the resulting DNA fragments were separated by agarose gel electrophoresis and then a fragment of 3.0 kb was recovered in the same manner.

Using a ligation kit, both of the fragments of 3.0 kb were subjected to ligation reaction for 16 hours at 16°C.

Using the ligation reaction solution, *Escherichia coli* NM522 was transformed in accordance with the usual process, and the resulting transformant was spread on LB agar medium containing 50 µg/ml ampicillin and then cultured overnight at 30°C.

By extracting a plasmid from the thus grown colonies of the transformant in accordance with the above-mentioned known process, plasmid pNK6 containing manC and manB was obtained. Structure of the plasmid was recognized by restriction enzyme cleavage (Fig. 11).

A 0.5 µg portion of the pNK6 DNA was cleaved with restriction enzymes ClaI and BamHI, the resulting DNA fragments were separated by agarose gel electrophoresis and then a fragment of 3.0 kb was recovered. A 0.2 µg portion of pPAC31 DNA obtained in Example 1-1) was cleaved with restriction enzymes ClaI and BamHI, the resulting DNA fragments were separated by agarose gel electrophoresis and then a fragment of 5.5 kb was recovered.

Using a ligation kit, the fragments of 3.0 kb and 5.5 kb were subjected to ligation reaction for 16 hours at 16°C.

Using the ligation reaction solution, *Escherichia coli* NM522 was transformed in accordance with the usual process, and the resulting transformant was spread on LB agar medium containing 50 µg/ml ampicillin and then cultured overnight at 30°C.

By extracting a plasmid from the thus grown colonies of the transformant in accordance with the usual process, *manC*, *manB* expression plasmid pNK7 was obtained. Structure of the plasmid was recognized by restriction enzyme cleavage (Fig. 11).

2) Construction of *pgm*, *pfkB* simultaneous expression plasmid

A 0.5 µg portion of the pNT24 DNA obtained in Example 7 was cleaved with restriction enzymes *Xho*I and *Bam*HI, the resulting DNA fragments were separated by agarose gel electrophoresis and then a fragment of 3.0 kb was recovered using Gene Clean II Kit. On the other hand, a 0.2 µg portion of pSTV28 DNA (manufactured by Takara Shuzo Co., Ltd.) was cleaved with restriction enzymes *Sal*I and *Bam*HI, the resulting DNA fragments were separated by agarose gel electrophoresis and then a fragment of 3.0 kb was recovered in the same manner.

Using a ligation kit, both of the fragments of 3.0 kb were subjected to ligation reaction for 16 hours at 16°C.

Using the ligation reaction solution, *Escherichia coli* NM522 was transformed in accordance with the usual process, and the resulting transformant was spread on LB agar medium containing 10 µg/ml chloramphenicol and then cultured overnight at 30°C.

By extracting a plasmid from the thus grown colonies of the transformant in accordance with the usual process, plasmid pNT53 containing the *pgm* gene was obtained. Structure of the plasmid was recognized by restriction enzyme cleavage (Fig. 12).

The sense strand DNA primer shown in SEQ ID NO:19 was synthesized, and the PCR was carried out using the sense strand DNA primer and the antisense strand DNA primer shown in SEQ ID NO:16, and the pNT47 DNA obtained in Example 7 as the template, under the same conditions as described in the foregoing.

After completion of the PCR, a precipitate of DNA was obtained by the ethanol precipitation process. The precipitate was dissolved in 20 µl of TE. Using 5 µl portion of the solution, the DNA was cleaved with restriction enzymes *EcoRV* and *BglII*, the resulting DNA fragments were separated by agarose gel electrophoresis and then a fragment of 1.3 kb was recovered in the same manner. A 0.2 µg portion of pNT53 DNA was cleaved with restriction enzymes *SmaI* and *BamHI*, the resulting DNA fragments were separated by agarose gel

electrophoresis and then a fragment of 6.0 kb was recovered in the same manner.

Using a ligation kit, the fragments of 1.3 kb and 6.0 kb were subjected to ligation reaction for 16 hours at 16°C.

Using the ligation reaction solution, *Escherichia coli* NM522 was transformed in accordance with the usual process, and the resulting transformant was spread on LB agar medium containing 10 µg/ml chloramphenicol and cultured overnight at 30°C.

By extracting a plasmid from the thus grown colonies of the transformant in accordance with the usual process, plasmid pNT55 capable of expressing *pgm* and *pfkB* was obtained. Structure of the plasmid was recognized by restriction enzyme cleavage (Fig. 12).

Example 14: Production of GDP-Man

1) Preparation of *manB*, *manC*, *pgm*, *pfkB* expression strain

Using the pNT55 DNA obtained in Example 13-2), *Escherichia coli* NM522/pNK7 was transformed in accordance with the usual process, and the resulting transformant was spread on LB agar medium containing 50 µg/ml ampicillin and 10 µg/ml chloramphenicol and cultured overnight at 30°C. By selecting the thus grown transformants, *Escherichia coli* NM522/pNK7/pNT55 was obtained as a *manB*, *manC*, *pgm*, *pfkB* expression strain.

2) Production of GDP-Man

Escherichia coli NM522/pNK7/pNT55 obtained in the above step 1) and *Escherichia coli* NM522/pNT46 obtained in Example 7 were cultured separately in the same manner as in Example 2, and each of thus obtained the culture broths was centrifuged to obtain wet cells. Also, *Corynebacterium ammoniagenes* ATCC 21170 was cultured in the same manner as in Example 2, and the culture broth was centrifuged to obtain wet cells. As occasion demands, these wet cells can be preserved at -20°C and utilized by thawing the cells prior to use.

A 30 ml portion of a reaction solution having a composition of 25 g/l *Escherichia coli* NM522/pNK7/pNT55 wet cells, 25 g/l the NM522/pNT46 wet cells, 150 g/l *Corynebacterium ammoniagenes* ATCC 21170 wet cells, 60 g/l fructose, 30 g/l mannose, 15 g/l KH_2PO_4 , 5 g/l $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 5 g/l phytic acid, 60 g/l GMP (2Na, $7\text{H}_2\text{O}$ salt), 4 g/l Nymean S-215 and 10 ml/l xylene was put into a 200 ml beaker, and 24 hours of the reaction was carried out under stirring the reaction solution with a magnetic stirrer (900 rpm).

During the reaction, pH of the reaction solution was maintained at 7.2 using 4 N NaOH, and KH_2PO_4 was added when required.

By the reaction, 14.6 g/l GDP-Man (2Na, $1\text{H}_2\text{O}$ salt) was formed in the reaction solution.

Example 15: Construction of *gmd*, *wcaG* expression plasmid

The sense strand DNA primer shown in SEQ ID NO:20 and the antisense strand DNA primer shown in SEQ ID NO:21 were synthesized. The PCR was carried out using the synthesized DNA strands as primers, and the chromosomal DNA of *Escherichia coli* W3110 as the template, under the same conditions as described in the foregoing.

After completion of the PCR, a precipitate of DNA was obtained by the ethanol precipitation process. The precipitate was dissolved in 20 μ l of TE. Using 5 μ l portion of the solution, the DNA was cleaved with restriction enzymes *Hind*III and *Xho*I, the resulting DNA fragments were separated by agarose gel electrophoresis and then a fragment of 2.3 kb was recovered using Gene Clean II Kit.

A 0.2 μ g portion of the pPA31 DNA obtained in Example 1-1) was cleaved with restriction enzymes *Hind*III and *Sal*I, the resulting DNA fragments were separated by agarose gel electrophoresis and then a fragment of 3.9 kb was recovered in the same manner.

Using a ligation kit, the fragments of 2.3 kb and 3.9 kb were subjected to ligation reaction for 16 hours at 16°C.

Using the ligation reaction solution, *Escherichia coli* NM522 was transformed in accordance with the usual process, and the resulting transformant was spread on LB agar

medium containing 50 µg/ml ampicillin and then cultured overnight at 30°C.

By extracting a plasmid from the thus grown colonies of the transformant in accordance with the usual process, plasmid pNK8 containing *gmd* and *wcaG* was obtained. Structure of the plasmid was recognized by restriction enzyme cleavage (Fig. 13).

Example 16: Production of GDP-Fuc

Escherichia coli NM522/pNK7/pNT55 obtained in Example 14, NM522/pNK8 obtained in Example 15 and NM522/pNT46 obtained in Example 7 were cultured in the same manner as in Example 2, and each of the thus obtained culture broths was centrifuged to obtain wet cells. Also, *Corynebacterium ammoniagenes* ATCC 21170 was cultured in the same manner as in Example 2, and the culture broth was centrifuged to obtain wet cells. As occasion demands, these wet cells can be preserved at -20°C and utilized by thawing the cells prior to use.

A 30 ml portion of a reaction solution having a composition of 25 g/l *Escherichia coli* NM522/pNK7/pNT55 wet cells, 25 g/l *Escherichia coli* NM522/pNK8 wet cells, 25 g/l *Escherichia coli* NM522/pNT46 wet cells, 150 g/l *Corynebacterium ammoniagenes* ATCC 21170 wet cells, 40 g/l fructose, 60 g/l mannose, 15 g/l KH_2PO_4 , 5 g/l $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 5

g/l phytic acid, 60 g/l GMP (2Na/7H₂O salt), 4 g/l Nymsean S-215 and 10 ml/l xylene was put into a 200 ml beaker, and 24 hours of the reaction was carried out at 32°C under stirring the reaction solution with a magnetic stirrer (900 rpm).

During the reaction, pH of the reaction solution was maintained at 7.2 using 4 N NaOH, and KH₂PO₄ was added when required.

By the reaction, 1.0 g/l GDP-Fuc (2.5Na, 1H₂O salt) was formed in the reaction solution.

Example 17: Construction of neuA expression plasmid

Chromosomal DNA of *Escherichia coli* K235 (ATCC 13027) was prepared in the same manner as in Example 1.

The sense strand DNA primer shown in SEQ ID NO:22 and the antisense strand DNA primer shown in SEQ ID NO:23 were synthesized. The PCR was carried out using the synthesized DNA strands as primers, and the chromosomal DNA of *Escherichia coli* K235 (ATCC 13027) as the template, under the same conditions as described in the foregoing.

After completion of the PCR treatment, a precipitate of DNA was obtained by the ethanol precipitation process. The precipitate was dissolved in 20 µl of TE. Using 5 µl portion of the solution, the DNA was cleaved with restriction enzymes EcoRI and BamHI, the resulting DNA fragments were separated by agarose gel electrophoresis and then a fragment

of 1.3 kb was recovered using Gene Clean II Kit. A 0.2 μ g portion of pBluescript II SK+ DNA was cleaved with restriction enzymes *EcoRI* and *BamHI*, the resulting DNA fragments were separated by agarose gel electrophoresis and then a fragment of 3.0 kb was recovered in the same manner.

Using a ligation kit, the fragments of 1.3 kb and 3.0 kb were subjected to ligation reaction for 16 hours at 16°C.

Using the ligation reaction solution, *Escherichia coli* NM522 was transformed in accordance with the usual process, and the resulting transformant was spread on LB agar medium containing 50 μ g/ml ampicillin and then cultured overnight at 30°C.

By extracting a plasmid from the thus grown colonies of the transformant in accordance with the usual process, plasmid pTA12 containing the *neuA* gene was obtained. Structure of the plasmid was recognized by restriction enzyme cleavage (Fig. 14).

A 0.5 μ g portion of the pTA12 DNA was cleaved with restriction enzymes *ClaI* and *BamHI*, the resulting DNA fragments were separated by agarose gel electrophoresis and then a fragment of 1.3 kb was recovered in the same manner. A 0.2 μ g portion of the pPAC31 DNA obtained in Example 1-1) was cleaved with restriction enzymes *ClaI* and *BamHI*, the resulting DNA fragments were separated by agarose gel

electrophoresis and then a fragment of 5.5 kb was recovered in the same manner.

Using a ligation kit, the fragments of 1.3 kb and 5.5 kb were subjected to ligation reaction for 16 hours at 16°C.

Using the ligation reaction solution, *Escherichia coli* NM522 was transformed in accordance with the usual process, and the resulting transformant was spread on LB agar medium containing 50 µg/ml ampicillin and then cultured overnight at 30°C.

By extracting a plasmid from the thus grown colonies of the transformant in accordance with the usual process, neuA expression plasmid pTA14 was obtained. Structure of the plasmid was recognized by restriction enzyme cleavage (Fig. 14).

Example 18: Production of CMP-NeuAc

Escherichia coli NM522/pTA14 obtained in Example 17, C600/pNAL1 (Appl. Environ. Microbiol., 51, 562 (1986)) and JF646/pM95 (J. Biol. Chem., 261, 5568 (1986)) were separately cultured in the same manner as in Example 2, and each of the thus obtained culture broths was centrifuged to obtain wet cells. Also, *Corynebacterium ammoniagenes* ATCC 21170 was cultured in the same manner as in Example 2, and the culture broth was centrifuged to obtain wet cells. As occasion

demands, these wet cells can be preserved at -20°C and utilized by thawing the cells prior to use.

A 30 ml portion of a reaction solution having a composition of 50 g/l *Escherichia coli* NM522/pTA14 wet cells, 15 g/l *Escherichia coli* C600/pNAL1 wet cells, 25 g/l *Escherichia coli* JF646/pMW5 wet cells, 150 g/l *Corynebacterium ammoniagenes* ATCC 21170 wet cells, 10 g/l orotic acid (potassium salt), 20 g/l pyruvic acid (Na salt), 40 g/l fructose, 10 g/l N-acetylmannosamine, 15 g/l KH_2PO_4 , 5 g/l $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 5 g/l phytic acid, 4 g/l Nymen S-215 and 10 ml/l xylene was put into a 200 ml beaker, and 24 hours of the reaction was carried out at 32°C under stirring the reaction solution with a magnetic stirrer (900 rpm).

During the reaction, pH of the reaction solution was maintained at 7.2 using 4 N NaOH, and KH_2PO_4 was added when required.

By the reaction, 2.7 g/l CMP-NeuAc (Na salt) was formed in the reaction solution.

Example 19: Production of lacto-N-tetraose

1) Preparation of β 1,3-galactosyltransferase

Namalwa cell line KJM-1 transformed with plasmid pAMoERSAW1 (Japanese Published Unexamined Patent Application No. 181759/94) containing a gene encoding a fusion protein of the IgG binding region of protein A with

Seikagaku Kogyo K.K.), and then the mixture was allowed to react for 16 hours at 37°C to remove the galactose at the non-reducing end.

The reaction solution was heated at 100°C for 5 minutes to inactivate β -galactosidase.

GlcNAc β 1-3Gal β 1-4Glc obtained by the reaction was used as a complex carbohydrate precursor.

A 36 μ l portion of a reaction solution containing 0.5 mM of the complex carbohydrate precursor, 0.5 U the β 1,3-galactosyltransferase linked IgG Sepharose obtained in the above step 1), 6 μ l of a reaction solution containing UDP-Gal (5 mM) obtained in Example 4, 100 mM Tris-HCl (pH 7.9), 10 mM MnCl₂, and 2 mM β -mercaptoethanol was allowed to stand for 65 hours at 32°C to effect the reaction.

After completion of the reaction, amount of the product accumulated in the reaction solution was measured by HPLC under the following conditions:

Column:

TSK gel ODS-80TM column (4.6 mm \times 30 cm, manufactured by TOSOH CORPORATION)

Liquid phase:

0.02 M ammonium acetate buffer (pH 4.0)

Temperature:

50°C

Flow rate:

1 ml/min

Detection:

Fluorescence detector (excitation wavelength 320 nm,
radiation wavelength 400 nm)

Identification of the product was carried out by
comparing elution time of aminopyridine-labeled lacto-N-
tetraose with that of the labeled product.

By the reaction, 0.17 mM (0.12 g/l) lacto-N-tetraose
was formed.

Example 20: Production of lacto-N-neotetraose

In the same manner as in Example 19, GlcNAc β 1-
3Gal β 1-4Glc was prepared from lacto-N-neotetraose and used as
a complex carbohydrate precursor.

A 36 μ l portion of a reaction solution containing 0.5
mM of the complex carbohydrate precursor, 0.5 U β 1,4-
galactosyltransferase (manufactured by Sigma), 6 μ l of a
reaction solution containing UDP-Gal (5 mM) obtained in
Example 4, 100 mM Tris-HCl buffer (pH 7.9), 10 mM MnCl₂ and 2
mM β -mercaptoethanol was allowed to stand for 65 hours at
32°C to effect the reaction.

After completion of the reaction, amount of the
product accumulated in the reaction solution was measured by
HPLC under the same conditions of Example 19-2).

Identification of the product was carried out by comparing elution time of the aminopyridine-labeled lacto-N-neotetraose with that of the product.

By the reaction, 0.15 mM (0.11 g/l) lacto-N-neotetraose was formed.

Example 21: Production of lacto-N-fucopentaose III

IgG Sepharose-linked α 1,3-fucosyltransferase was prepared from namalwa cell line KJM-1 transformed with plasmid pAMCA-FT6 (*J. Biol. Chem.*, 269, 14730 (1994)) containing a gene encoding a fusion protein of the IgG binding region of protein A with α 1,3-fucosyltransferase, in the same manner as in Example 19-1), and used as the enzyme source of α 1,3-fucosyltransferase.

A 50 μ l portion of a reaction solution containing 0.25 mM lacto-N-neotetraose (manufactured by Oxford Glycosystems), 1.0 U of the IgG Sepharose-linked α 1,3-fucosyltransferase, 6 μ l of a reaction solution containing GDP-Fuc (0.25 mM) obtained in Example 16, 100 mM Tris-HCl buffer (pH 7.9) and 10 mM MnCl₂ was allowed to stand for 24 hours at 37°C to effect the reaction.

After completion of the reaction, amount of the product accumulated in the reaction solution was measured using a sugar analyzer (DX-500) manufactured by Dionex. Identification of the product was carried out by comparing

elution time of lacto-N-fucopentaose III (manufactured by Oxford Glycosystems) with that of the product.

By the reaction, 0.21 mM (0.18 g/l) lacto-N-fucopentaose III was formed.

Example 22: Construction of α 1,4-galactosyltransferase (lgtC) expression plasmid

Chromosomal DNA of *Neisseria gonorrhoeae* (ATCC 33084) was prepared in the same manner as in Example 1.

The sense strand DNA primer shown in SEQ ID NO:24 and the antisense strand DNA primer shown in SEQ ID NO:25 were synthesized. The PCR was carried out using the synthesized DNA strands as primers, and the chromosomal DNA of *Neisseria gonorrhoeae* (ATCC 33084) as the template, under the same conditions as described in the foregoing.

After completion of the PCR treatment, a precipitate of DNA was obtained by the ethanol precipitation process. The precipitate was dissolved in 20 μ l of TE. Using 5 μ l portion of the solution, the DNA was cleaved with restriction enzymes *Hind*III and *Bam*HI, the resulting DNA fragments were separated by agarose gel electrophoresis and then a fragment of 1.0 kb was recovered using Gene Clean II Kit. A 0.2 μ g portion of the pPA31 DNA obtained in Example 1-1) was cleaved with restriction enzymes *Hind*III and *Bam*HI, the resulting DNA

fragments were separated by agarose gel electrophoresis and then a fragment of 4.2 kb was recovered in the same manner.

Using a ligation kit, the fragments of 1.0 kb and 4.2 kb were subjected to ligation reaction for 16 hours at 16°C.

Using the ligation reaction solution, *Escherichia coli* NM522 was transformed in accordance with the usual process, and the resulting transformant was spread on LB agar medium containing 50 µg/ml ampicillin and then cultured overnight at 30°C.

By extracting a plasmid from the thus grown colonies of the transformant in accordance with the usual process, a lgtC expression plasmid pGT3 was obtained. Structure of the plasmid was recognized by restriction enzyme cleavage (Fig. 15).

Example 23: Production of globotriose

Escherichia coli NM522/pNT25/pNT32 obtained in Example 4 and *Escherichia coli* NM522/pGT3 obtained in Example 22 were cultured in the same manner as in Example 2, and each of the thus obtained culture broths was centrifuged to obtain wet cells. Also, *Corynebacterium ammoniagenes* ATCC 21170 was cultured in the same manner as in Example 2, and the culture broth was centrifuged to obtain wet cells. As occasion demands, these wet cells can be preserved at -20°C and utilized by thawing the cells prior to use.

[illegible]

項目	単位	数値	単位	数値	単位	数値	単位	数値
1. 総人口	人	1,234,567	2. 男性人口	人	612,345	3. 女性人口	人	622,222
4. 0歳人口	人	12,345	5. 1歳人口	人	11,234	6. 2歳人口	人	10,123
7. 3歳人口	人	9,012	8. 4歳人口	人	8,901	9. 5歳人口	人	7,890
10. 6歳人口	人	6,789	11. 7歳人口	人	5,678	12. 8歳人口	人	4,567
13. 9歳人口	人	3,456	14. 10歳人口	人	2,345	15. 11歳人口	人	1,234
16. 12歳人口	人	1,123	17. 13歳人口	人	1,012	18. 14歳人口	人	901
19. 15歳人口	人	890	20. 16歳人口	人	789	21. 17歳人口	人	678
22. 18歳人口	人	567	23. 19歳人口	人	456	24. 20歳人口	人	345
25. 21歳人口	人	234	26. 22歳人口	人	123	27. 23歳人口	人	112
28. 24歳人口	人	101	29. 25歳人口	人	90	30. 26歳人口	人	89
31. 27歳人口	人	78	32. 28歳人口	人	67	33. 29歳人口	人	56
34. 30歳人口	人	45	35. 31歳人口	人	34	36. 32歳人口	人	23
37. 33歳人口	人	12	38. 34歳人口	人	11	39. 35歳人口	人	10
40. 36歳人口	人	9	41. 37歳人口	人	8	42. 38歳人口	人	7
43. 39歳人口	人	6	44. 40歳人口	人	5	45. 41歳人口	人	4
46. 42歳人口	人	3	47. 43歳人口	人	2	48. 44歳人口	人	1
49. 45歳人口	人	1	50. 46歳人口	人	0	51. 47歳人口	人	0
52. 48歳人口	人	0	53. 49歳人口	人	0	54. 50歳人口	人	0
55. 51歳人口	人	0	56. 52歳人口	人	0	57. 53歳人口	人	0
58. 54歳人口	人	0	59. 55歳人口	人	0	60. 56歳人口	人	0
61. 57歳人口	人	0	62. 58歳人口	人	0	63. 59歳人口	人	0
64. 60歳人口	人	0	65. 61歳人口	人	0	66. 62歳人口	人	0
67. 63歳人口	人	0	68. 64歳人口	人	0	69. 65歳人口	人	0
70. 66歳人口	人	0	71. 67歳人口	人	0	72. 68歳人口	人	0
73. 69歳人口	人	0	74. 70歳人口	人	0	75. 71歳人口	人	0
76. 72歳人口	人	0	77. 73歳人口	人	0	78. 74歳人口	人	0
79. 75歳人口	人	0	80. 76歳人口	人	0	81. 77歳人口	人	0
82. 78歳人口	人	0	83. 79歳人口	人	0	84. 80歳人口	人	0
85. 81歳人口	人	0	86. 82歳人口	人	0	87. 83歳人口	人	0
88. 84歳人口	人	0	89. 85歳人口	人	0	90. 86歳人口	人	0
91. 87歳人口	人	0	92. 88歳人口	人	0	93. 89歳人口	人	0
94. 90歳人口	人	0	95. 91歳人口	人	0	96. 92歳人口	人	0
97. 93歳人口	人	0	98. 94歳人口	人	0	99. 95歳人口	人	0
100. 96歳人口	人	0	101. 97歳人口	人	0	102. 98歳人口	人	0
103. 99歳人口	人	0	104. 100歳人口	人	0	105. 101歳人口	人	0
106. 102歳人口	人	0	107. 103歳人口	人	0	108. 104歳人口	人	0
109. 105歳人口	人	0	110. 106歳人口	人	0	111. 107歳人口	人	0
112. 108歳人口	人	0	113. 109歳人口	人	0	114. 110歳人口	人	0
115. 111歳人口	人	0	116. 112歳人口	人	0	117. 113歳人口	人	0
118. 114歳人口	人	0	119. 115歳人口	人	0	120. 116歳人口	人	0
121. 117歳人口	人	0	122. 118歳人口	人	0	123. 119歳人口	人	0
124. 120歳人口	人	0	125. 121歳人口	人	0	126. 122歳人口	人	0
127. 123歳人口	人	0	128. 124歳人口					

Figure 1

600

each of the thus obtained culture broths was centrifuged to obtain wet cells. Also, *Corynebacterium ammoniagenes* ATCC 21170 was cultured in the same manner as in Example 2, and the culture broth was centrifuged to obtain wet cells. As occasion demands, these wet cells can be preserved at -20°C and utilized by thawing the cells prior to use.

A 30 ml portion of a reaction solution having a composition of 50 g/l *Escherichia coli* NM522/pNT25/pNT32 wet cells, 50 g/l *Escherichia coli* NM522/pGT3 wet cells, 150 g/l *Corynebacterium ammoniagenes* ATCC 21170 wet cells, 50 g/l fructose, 50 g/l galactose, 96 g/l N-acetyllactosamine, 15 g/l KH_2PO_4 , 5 g/l $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 5 g/l phytic acid, 10 g/l orotic acid (potassium salt), 4 g/l Nymean S-215 and 10 ml/l xylene was put into a 200 ml beaker, and 23 hours of the reaction was carried out at 32°C under stirring the reaction solution with a magnetic stirrer (900 rpm).

During the reaction, pH of the reaction solution was maintained at 7.2 using 4 N NaOH, and galactose, fructose and KH_2PO_4 were added when required.

By the reaction, 10 g/l Gal α 1-4Gal β 1-4GlcNAc was formed in the reaction solution.

Cells were removed from the reaction solution by centrifugation, and the formed product was purified from a 30 ml portion of the thus obtained supernatant by employing a

process in which activated carbon was used, thereby obtaining 0.2 g of Gal α 1-4Gal β 1-4GlcNAc as white powder.

Example 25: Construction of β 1,4-galactosyltransferase (lgTB) expression plasmid

The sense strand DNA primer shown in SEQ ID NO:26 and the antisense strand DNA primer shown in SEQ ID NO:27 were synthesized. The PCR was carried out using the synthesized DNA strands as primers, and the chromosomal DNA of *N. gonorrhoeae* (ATCC 33084) as the template, under the same conditions as described in the foregoing.

After completion of the PCR treatment, a precipitate of DNA was obtained by the ethanol precipitation process. The precipitate was dissolved in 20 μ l of TE. Using 5 μ l portion of the solution, the DNA was cleaved with restriction enzymes *Hind*III and *Bam*HI, the resulting DNA fragments were separated by agarose gel electrophoresis and then a fragment of 0.8 kb was recovered using Gene Clean II Kit. A 0.2 μ g portion of pBluescript II SK+ DNA was cleaved with restriction enzymes *Hind*III and *Bam*HI, the resulting DNA fragments were separated by agarose gel electrophoresis and then a fragment of 3.0 kb was recovered in the same manner.

Using a ligation kit, the fragments of 0.8 kb and 3.0 kb were subjected to ligation reaction for 16 hours at 16°C.

Using the ligation reaction solution, *Escherichia coli* NM522 was transformed in accordance with the usual process, and the resulting transformant was spread on LB agar medium containing 50 µg/ml ampicillin and then cultured overnight at 30°C.

By extracting a plasmid from the thus grown colonies of the transformant in accordance with the usual process, plasmid pNT60P containing the lgtB gene was obtained. Structure of the plasmid was recognized by restriction enzyme cleavage (Fig. 16).

A 0.5 µg portion of the pNT60P DNA was cleaved with restriction enzymes *Cla*I and *Bam*HI, the resulting DNA fragments were separated by agarose gel electrophoresis and then a fragment of 0.8 kb was recovered. A 0.2 µg portion of the pPAC31 DNA obtained in Example 1-1) was cleaved with restriction enzymes *Cla*I and *Bam*HI, the resulting DNA fragments were separated by agarose gel electrophoresis and then a fragment of 5.5 kb was recovered in the same manner.

Using a ligation kit, the fragments of 0.8 kb and 5.5 kb were subjected to ligation reaction for 16 hours at 16°C.

Using the ligation reaction solution, *Escherichia coli* NM522 was transformed in accordance with the usual process, and the resulting transformant was spread on LB agar medium containing 50 µg/ml ampicillin and then cultured overnight at 30°C.

By extracting a plasmid from the thus grown colonies of the transformant in accordance with the usual process, a lgtB expression plasmid pNT60 was obtained. Structure of the plasmid was recognized by restriction enzyme cleavage (Fig. 16).

Example 26: Production of N-acetyllactosamine

Escherichia coli NMS22/pNT60 obtained in Example 25 and *Escherichia coli* NMS22/pNT25 obtained in Example 3 were cultured in the same manner as in Example 2, and each of the thus obtained culture broths was centrifuged to obtain wet cells. Also, *Corynebacterium ammoniagenes* ATCC 21170 was cultured in the same manner as in Example 2, and the culture broth was centrifuged to obtain wet cells. As occasion demands, these wet cells can be preserved at -20°C and utilized by thawing the cells prior to use.

A 30 ml portion of a reaction solution having a composition of 50 g/l *Escherichia coli* NMS22/pNT25 wet cells, 50 g/l *Escherichia coli* NMS22/pNT60 wet cells, 150 g/l *Corynebacterium ammoniagenes* ATCC 21170 wet cells, 10 g/l orotic acid (potassium salt), 100 g/l fructose, 100 g/l N-acetylglucosamine, 100 g/l galactose, 15 g/l KH_2PO_4 , 5 g/l $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 5 g/l phytic acid, 4 g/l Nymeen S-215 and 10 ml/l xylene was put into a 200 ml beaker, and 34 hours of the

reaction was carried out at 32°C under stirring the reaction solution with a magnetic stirrer (900 rpm).

During the reaction, pH of the reaction solution was maintained at 7.2 using 4 N NaOH, and galactose, fructose and KH_2PO_4 were added when required.

By the reaction, 114 g/l N-acetyllactosamine was formed in the reaction solution.

Example 27: Production of lactose

Escherichia coli NM522/pNT60 obtained in Example 25 and *Escherichia coli* NM522/pNT25 obtained in Example 3 were cultured in the same manner as in Example 2, and each of the thus obtained culture broths was centrifuged to obtain wet cells. Also, *Corynebacterium ammoniagenes* ATCC 21170 was cultured in the same manner as in Example 2, and the culture broth was centrifuged to obtain wet cells. As occasion demands, these wet cells can be preserved at -20°C and utilized by thawing the cells prior to use.

A 30 ml portion of a reaction solution having a composition of 50 g/l *Escherichia coli* NM522/pNT25 wet cells, 50 g/l *Escherichia coli* NM522/pNT60 wet cells, 150 g/l *Corynebacterium ammoniagenes* ATCC 21170 wet cells, 10 g/l orotic acid (potassium salt), 115 g/l glucose, 115 g/l galactose, 15 g/l KH_2PO_4 , 5 g/l $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 5 g/l phytic acid, 4 g/l Nymean S-215 and 10 ml/l xylene was put into a 200 ml

beaker, and 15 hours of the reaction was carried out at 32°C by stirring the reaction solution on a magnetic stirrer (900 rpm).

During the reaction, pH of the reaction solution was maintained at 7.2 using 4 N NaOH, and KH_2PO_4 was added when required.

By the reaction, 49 g/l lactose was formed in the reaction solution.

Example 28: Production of globotriose

Escherichia coli NM522/pNT60 obtained in Example 25, *Escherichia coli* NM522/pNT25 obtained in Example 3 and *Escherichia coli* NM522/pGT3 obtained in Example 22 were cultured in the same manner as in Example 2, and each of the thus obtained culture broths was centrifuged to obtain wet cells. Also, *Corynebacterium ammoniagenes* ATCC 21170 was cultured in the same manner as in Example 2, and the culture broth was centrifuged to obtain wet cells. As occasion demands, these wet cells can be preserved at -20°C and utilized by thawing the cells prior to use.

A 30 ml portion of a reaction solution having a composition of 50 g/l *Escherichia coli* NM522/pNT25 wet cells, 50 g/l *Escherichia coli* NM522/pNT60 wet cells, 50 g/l *Escherichia coli* NM522/pGT3 wet cells, 150 g/l *Corynebacterium ammoniagenes* ATCC 21170 wet cells, 10 g/l

orotic acid (potassium salt), 115 g/l glucose, 115 g/l galactose, 15 g/l KH_2PO_4 , 5 g/l $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 5 g/l phytic acid, 4 g/l Nymean S-215 and 10 ml/l xylene was put into a 200 ml beaker, and 13 hours of the reaction was carried out at 32°C under stirring the reaction solution with a magnetic stirrer (900 rpm).

During the reaction, pH of the reaction solution was maintained at 7.2 using 4 N NaOH, and KH_2PO_4 was added when required.

By the reaction, 5 g/l globotriose was formed in the reaction solution.

INDUSTRIAL APPLICABILITY

The present invention renders possible efficient industrial production of a sugar nucleotide from a nucleotide precursor and a sugar as the sole starting materials and of a complex carbohydrate from the sugar nucleotide and a complex carbohydrate precursor.